



# Three Year Trending & Benchmark Analysis

## Molina Healthcare of New Mexico

Medicaid	HEDIS Reporting Year			Rate Change*		Medicaid Plans Accreditation Thresholds 2008				NCOA Medicaid Benchmarks -- 2007					
	2006	2007	2008	2006 to 2008	2007 to 2008	25th	50th	75th	90th	Molina Ranking**		50th	75th	90th	Molina Ranking
<b>Childhood Immunization Status</b>															
DT/DTP/DTPaP	71.02%	83.33%	81.88%	10.86%	-1.45%							79	84.1	88.9	Above 50TH
OPV/IPV	76.99%	90.74%	91.06%	14.07%	0.32%							87.8	91.7	94.7	Above 50TH
Measles, Mumps, & Rubella	81.64%	93.06%	94.44%	12.80%	1.38%							90.8	93.4	95.3	Above 75TH
H Influenza Type B	78.98%	90.51%	90.34%	11.36%	-0.17%							89.5	92.4	95.1	Above 50TH
Hepatitis B	71.90%	87.04%	90.58%	18.68%	3.54%							88	92.2	95.2	Above 50TH
Chicken Pox	79.20%	91.44%	91.30%	12.10%	-0.14%							88.6	92	93.8	Above 50TH
Pneumococcal Conjugate	28.98%	74.31%	78.26%	49.28%	3.95%							47.3	55	64.2	Above 90TH
Combination 2	56.64%	74.07%	76.09%	19.45%	2.02%	63	72	79	83	Above 90TH		72.4	78.5	82.7	Above 50TH
Combination 3	23.67%	65.28%	71.01%	47.34%	5.73%							42.3	49.8	57.8	Above 90TH
<b>Well Child Care: First 15 months</b>															
Zero visits	7.73%	2.08%	0.93%	-6.80%	-1.15%							1.4	2.9	6.8	
One visit	6.40%	1.16%	0.93%	-5.47%	-0.23%							1.7	3.1	5.1	
Two visits	7.06%	3.47%	1.16%	-5.90%	-2.31%							3.2	4.7	6.5	
Three visits	8.39%	7.87%	5.32%	-3.07%	-2.55%							5.8	7.9	9.5	
Four visits	16.11%	15.74%	11.81%	-4.30%	-3.93%							10.5	14.1	16.3	Above 50TH
Five visits	18.98%	19.68%	17.13%	-1.85%	-2.55%							17.4	21	23.8	
Six or more visits	35.32%	50.00%	62.73%	27.41%	12.73%							56.6	64.4	75.2	Above 50TH
<b>Well Child Care: Years 3-6</b>	<b>67.33%</b>	<b>61.81%</b>	<b>58.74%</b>	<b>-8.59%</b>	<b>-3.07%</b>							<b>67.5</b>	<b>74.9</b>	<b>79.9</b>	
<b>Well Child Care: Adolescence</b>	<b>43.49%</b>	<b>41.20%</b>	<b>50.93%</b>	<b>7.44%</b>	<b>9.73%</b>							<b>42.1</b>	<b>51.4</b>	<b>58.9</b>	<b>Above 50TH</b>
<b>Appropriate Testing for Pharyngitis</b>	<b>45.30%</b>	<b>47.93%</b>	<b>50.95%</b>	<b>5.65%</b>	<b>3.02%</b>	<b>47</b>	<b>59</b>	<b>69</b>	<b>76</b>	<b>Above 50TH</b>		<b>59.6</b>	<b>69.1</b>	<b>76.9</b>	
<b>URI Treatment</b>	<b>79.23%</b>	<b>86.52%</b>	<b>81.86%</b>	<b>2.63%</b>	<b>-4.66%</b>	<b>78</b>	<b>84</b>	<b>89</b>	<b>93</b>	<b>Above 90TH</b>		<b>84.3</b>	<b>89.4</b>	<b>92.6</b>	
<b>Breast Cancer Screening</b>	<b>57.63%</b>	<b>56.41%</b>	<b>51.95%</b>	<b>-5.68%</b>	<b>-4.46%</b>	<b>47</b>	<b>54</b>	<b>59</b>	<b>65</b>	<b>Above 75TH</b>		<b>54.9</b>	<b>59.2</b>	<b>65.2</b>	
<b>Cervical Cancer Screening</b>	<b>69.56%</b>	<b>64.21%</b>	<b>67.48%</b>	<b>-2.08%</b>	<b>3.27%</b>	<b>49</b>	<b>60</b>	<b>74</b>	<b>79</b>	<b>Above 90TH</b>		<b>66.5</b>	<b>72.0</b>	<b>77.4</b>	<b>Above 50TH</b>
<b>Chlamydia Screening in Women</b>															
Ages 16 to 20	36.66%	44.74%	48.85%	12.19%	4.11%							50.3	58.6	64.8	
Ages 21 to 25	35.86%	49.57%	52.59%	16.73%	3.02%							56.3	63	69.9	
Total	36.28%	46.94%	50.43%	14.15%	3.49%							52.8	60.6	66	
<b>Prenatal/Postpartum Care<sup>R</sup></b>															
Timeliness of prenatal care	89.28%	76.34%	76.34%	-12.94%	0.00%	74.0	81.0	86.0	90.0	Above 50TH		84.2	88.7	91.5	
Postpartum care	56.41%	49.78%	49.78%	-6.63%	0.00%	45	55	62	67	Above 50TH		59.7	65.5	71.1	
<b>Frequency of Prenatal Care<sup>R</sup></b>															
Less than 21 percent	13.24%	18.75%	18.75%	5.51%	0.00%							7.6	17.5	32.0	Above 75TH
21 to 40 percent	8.27%	10.04%	10.04%	1.77%	0.00%							4.6	7.8	12.9	Above 75TH
41 to 60 percent	7.80%	9.38%	9.38%	1.58%	0.00%							7.4	9.5	13.2	Above 50TH
61 to 80 percent	14.42%	16.96%	16.96%	2.54%	0.00%							13.9	17.0	20.6	Above 50TH
More than 80 percent	56.26%	44.87%	44.87%	-11.39%	0.00%							62.9	71.7	78.6	
<b>Controlling High Blood Pressure<sup>R</sup></b>															
	57.55%	57.74%	57.74%	0.19%	0.00%	47	56	60	66	Above 90TH		55.4	59.9	65.8	Above 50TH
<b>Cholesterol Management</b>															
LDL Screening	64.82%	80.34%	80.23%	15.41%	-0.11%	71	78	82	87	Above 90TH					
LDL < 100	23.27%	36.52%	40.68%	17.41%	4.16%										
<b>Comprehensive Diabetes Care</b>															
HbA1c Testing	84.80%	86.58%	79.88%	-4.92%	-6.70%	70	79	84	89	Above 75TH		79.3	84.3	89.1	Above 50TH
HbA1c Control	NA	28.72%	25.81%	NA	-2.91%							31.3	36.6	40.9	
HbA1c Poor Control	45.18%	51.78%	54.67%	9.49%	2.89%	58	48	38	31			46.7	39.7	32.1	
Eye Exam	60.69%	61.01%	55.49%	-5.20%	-5.52%	42	54	63	68	Above 90TH		53.6	62.7	68.3	Above 50TH
LDL-C Screening	83.30%	76.31%	70.93%	-12.37%	-5.38%	67	73	78	81	Above 50TH		72.8	77.9	81	
LDL Less than 100	37.90%	32.70%	29.07%	-8.83%	-3.63%							31.3	37.2	44.1	
Monitoring for Nephropathy	50.32%	79.66%	74.59%	24.27%	-5.07%	69	77	82	86	Above 50TH		76.6	81.8	85.5	

\*Shading indicates significance according to current NCOA accreditation guidelines

\*\*Using 5 point and appropriate regional adjustments

<sup>R</sup>Retired indicator <sup>R</sup>Rotated measure



## Three Year Trending & Benchmark Analysis

### Molina Healthcare of New Mexico

Medicaid	HEDIS Reporting Year			Rate Change*		Medicaid Plans Accreditation Thresholds 2008				NCOA Medicaid Benchmarks -- 2007					
	2006	2007	2008	2006 to 2008	2007 to 2008	25th	50th	75th	90th	Molina Ranking**		50th	75th	90th	Molina Ranking
<b>Appropriate Medication for Asthma</b>															
Ages 5 to 9	89.74%	94.00%	92.11%	2.37%	-1.89%							91.7	94.6	96.3	
Ages 10 to 17	85.71%	87.12%	84.59%	-1.12%	-2.53%							88.8	91.4	93	
Ages 18 to 56	79.28%	95.08%	80.55%	1.27%	-14.53%							85.4	88.2	90.9	
Combined Rate	84.52%	91.81%	85.31%	0.79%	-6.50%	84	87	90	93	Above 90TH		88.4	90.3	92	
<b>Antidepressant Medication Management</b>															
Optimal Practitioner Contacts	NA	NA	NA	NA	NA	12	23	37	44			22.0	27.0	31.8	
Effective Acute Phase Treatment	NA	NA	NA	NA	NA	40	45	52	65			44.7	47.9	51.1	
Effective Continuation Phase Treatment	NA	NA	NA	NA	NA	22	32	38	46			28.5	32.3	34.8	
<b>Follow-up Visit after MH Hospitalization</b>															
Follow Up within 30 Days	NA	NA	NA	NA	NA							57.3	75.9	79.8	
Follow Up within 7 Days	NA	NA	NA	NA	NA	26	38	58	66			35.8	58.0	66.2	
<b>Initiation/Engagement of AOD Services</b>															
Initiation - Total	43.58%	NA	NA	NA	NA							41.5	50.0	57.5	
Engagement - Total	9.44%	NA	NA	NA	NA							9.4	15.1	23.6	
<b>Childrens Access to Primary Care to</b>															
Ages 12-24 mos.	97.79%	97.82%	97.64%	-0.15%	-0.18%							95.8	97.4	98.0	Above 75TH
Ages 25 mos to 6 yrs.	85.75%	89.14%	89.25%	3.50%	0.11%							86.7	89.4	91.4	Above 50TH
Ages 7-11 yrs.	74.63%	90.74%	92.61%	17.98%	1.87%							87.2	90.5	93.3	Above 75TH
Ages 12-19 yrs.	73.39%	89.98%	91.71%	18.32%	1.73%							85.3	89.2	91.4	Above 90TH
<b>Adult Access to Preventive Care</b>															
Ages 20-44	84.32%	83.55%	84.38%	0.06%	0.83%							79.1	85.1	88.0	Above 50TH
Ages 45-64	87.47%	88.78%	88.34%	0.87%	-0.44%							85.5	88.6	89.8	Above 50TH
Ages 65 and older	87.07%	81.88%	82.38%	-4.69%	0.50%							82.3	88.7	93.5	Above 50TH
<b>Annual Dental Visit</b>															
Ages 2-3	25.90%	41.89%	46.41%	20.51%	4.52%										
Ages 4-6	53.15%	69.26%	71.04%	17.89%	1.78%							51.2	61.5	65.7	Above 90TH
Ages 7-10	63.34%	76.91%	77.30%	13.96%	0.39%							52.7	58.0	70.2	Above 90TH
Ages 11-14	57.87%	70.09%	71.90%	14.03%	1.81%							46.4	53.8	61.0	Above 90TH
Ages 15-18	45.96%	58.00%	58.30%	12.34%	0.30%							38.3	47.8	52.8	Above 90TH
Ages 19-21	32.94%	38.90%	40.79%	7.85%	1.89%							30.6	36.9	42.5	Above 75TH
Combined	50.86%	64.62%	66.01%	15.15%	1.39%							42.8	50.6	57.3	Above 90TH
<b>Advice to Quit Smoking</b>	63.68%	59.92%	63.82%	0.14%	3.90%	65	68	72	76	Above 50TH					
<b>Avoidance of Antibiotic Treatment</b>	24.18%	18.53%	12.32%	-11.86%	-6.21%	21.0	25.0	29.0	34.0			24.6	30.2	40.7	
<b>Imaging Studies for Low Back Pain</b>	72.55%	76.45%	78.21%	5.66%	1.76%	75.0	78.0	83.0	85.0			78.2	82.5	85.2	Above 50TH
<b>Persistence of Beta-Blocker Tx</b>	NA	NA	NA	NA	NA	62.0	72.0	78.0	81.0						

\*Shading indicates significance according to current NCOA accreditation guidelines

\*\*Using 5 point and appropriate regional adjustments

\*Retired indicator <sup>R</sup>Rotated measure