

# Immunization Schedule

(Birth to Six months old)

Caring For Your New Baby

## Do you have questions?

Call our 24-hour Nurse Advice Line.

We are here to help you.



English:

**(888) 275-8750**

Español:

**(866) 648-3537**

TTY/TDD:

**711**

Distributed by Molina Healthcare.  
All material in this brochure is for  
information only. It does not replace  
your provider's advice.

To get this information in other languages  
and accessible formats, please call Member  
Services. This number is on the back of  
your Member ID card.



[MolinaHealthcare.com](http://MolinaHealthcare.com)



Your Extended Family



Your Extended Family

## Keep your baby's appointments.

This is key to ensure your baby gets the right shots at the right time.

|                           |                                | BIRTH | 1 MO.             | 2 MO. | 4 MO. | 6 MO.  |
|---------------------------|--------------------------------|-------|-------------------|-------|-------|--|
| RECOMMENDED IMMUNIZATIONS | HEPATITIS B                    | HepB  | HepB <sup>1</sup> |       | —     | HepB   |
|                           | ROTAVIRUS                      | —     | —                 | RV    | RV    | RV   |
|                           | DIPHTHERIA, TETANUS, PERTUSSIS | —     | —                 | DTaP  | DTaP  | DTaP   |
|                           | HAEMOPHILUS INFLUENZAE TYPE B  | —     | —                 | Hib   | Hib   | Hib  |
|                           | PNEUMOCOCCAL                   | —     | —                 | PCV   | PCV   | PCV  |
|                           | INACTIVATED POLIOVIRUS         | —     | —                 | IPV   | IPV   | IPV  |
|                           | INFLUENZA                      | —     | —                 | —     | —     | Influenza<br>1 <sup>st</sup> dose <sup>2</sup><br>2 <sup>nd</sup> dose |

<sup>1</sup> The second dose of HepB may be given either at the one month or two month visit.

<sup>2</sup> Two doses of Influenza, given at least four weeks apart, are recommended for children aged six months through eight years of age who are getting a flu vaccine for the first time.



U.S. Department of Health and Human Services  
Centers for Disease Control and Prevention



Disease spreads easily from child to child. Immunization shots **protect your baby** from harmful disease and infection.

The shots are **safe** and they work!

**Keep track** of your baby's shots. Your baby's provider should remind you when your baby is due for his or her next shot.