



# Immunization Timing

Age 2 months	Interval from previous dose
DTaP	
Hib	
PCV (Pneumo)	
Polio (IPV)	
HepB	1-2 months after birth dose
RV (Rotavirus)	4-10 weeks

Age 4 months	Interval from previous dose
DTaP	1-2 months
Hib	1-2 months
PCV (Pneumo)	1-2 months
Polio (IPV)	1-2 months
HepB	1-2 months if 1st dose given at 2 months of age
RV (Rotavirus)	4-10 weeks

Age 6 months	Interval from previous dose
DTaP	1-2 months
Hib	1-2 months
PCV (Pneumo)	1-2 months
Polio (IPV)	1-14 months
HepB	2-12 months and at least 4 months after first dose
RV (Rotavirus)	4-10 weeks

Age 12 months	Interval from previous dose
DTaP	6-12 months
Hib	2-8 months
PCV (Pneumo)	6-8 weeks
MMR	
Varicella	
HepA	

Age 18 months	Interval from previous dose
HepA	6-12 months

**For School-Age Kids**

**Before Kindergarten**

Polio, DTaP, MMR #2

**Before 7<sup>th</sup> Grade**

Varicella,  
Hep B  
Tdap booster

Every Fall: 6-59 months and high-risk kids  
Flu Vaccine

This is a suggested schedule. For alternatives and details, consult the latest "Recommended Childhood Immunization Schedule, United States 2006."