

Non-HEDIS Measure Tips:

Tobacco Cessation Counseling

MEASURE DESCRIPTION

Patients 12 years of age and older (for BC+) and 19 years of age and older (for SSI) who were diagnosed as tobacco users in the calendar year and who received tobacco cessation counseling during the calendar year. Counseling to quit can be either face-to-face or by phone as identified by any claim or encounter with at least one of the codes for counseling and one of the tobacco user diagnosis codes, listed in the tables below, in the same encounter. Members must have continuous enrollment during the measurement year with no more than a one month gap in coverage. This is a non-HEDIS measure developed by DHS.

USING CORRECT BILLING CODES

Codes to Identify Tobacco Cessation Counseling:

Description	Codes
Tobacco User Diagnosis:	ICD-10-CM: F17200, F17201, F17203, F17208, F17209, F17210, F17211, F17213, F17218, F17219, F17220, F17221, F17223, F17228, F17229, F17290, F17291, F17293, F17298, F17299, Z720

Description	Codes
Codes to Identify Tobacco Cessation Counseling	CPT: 96150-96154, 99201-99205, 99211-99215, 99241-99245, 99384-99387, 99394-99397, 99401 -99404, 99406, 99407, 90832-90834, 90836-90838, 90845, 90847, 90849, 90853, 90875, 90876, 90880, 90899, 98967-98968, 99442-99443
	HCPCS: G9016 S9453

Description	Codes	
Exclusions:	History of Tobacco Use Diagnosis	ICD-10-CM: Z87891
	Pregnancy Diagnosis	ICD-10 CM: O0000 – O9A53, Z331-Z333, Z3400-Z3493, Z360-Z369, Z3A00-Z3A49, Z370-Z379
	Tobacco Use Disorder Complicating Pregnancy	ICD-10-CM: O99330, O99331, O99332, O99333