

Recommendations for Preventive Pediatric Health Care

Bright Futures/American Academy of Pediatrics

Each child and family is unique; therefore, these Recommendations for Preventive Pediatric Health Care are designed for the care of children who are receiving competent parenting, have no manifestations of any important health problems, and are growing and developing in satisfactory fashion. Additional visits may become necessary if circumstances suggest variations from normal.

Developmental, psychosocial, and chronic disease issues for children and adolescents may require frequent counseling and treatment visits separate from preventive care visits.

These guidelines represent a consensus by the American Academy of Pediatrics (AAP) and Bright Futures. The AAP continues to emphasize the great importance of continuity of care in comprehensive health supervision and the need to avoid fragmentation of care.

Refer to the specific guidance by age as listed in *Bright Futures* guidelines (Hagan JF, Shaw JS, Duncan PM, eds. *Bright Futures Guidelines for Health Supervision of Infants, Children and Adolescents*. 3rd ed. Elk Grove Village, IL: American Academy of Pediatrics; 2008).

The recommendations in this statement do not indicate an exclusive course of treatment or standard of medical care. Variations, taking into account individual circumstances, may be appropriate.

Copyright © 2016 by the American Academy of Pediatrics, updated 10/2015.

No part of this statement may be reproduced in any form or by any means without prior written permission from the American Academy of Pediatrics except for one copy for personal use.

AGE ¹	INFANCY								EARLY CHILDHOOD						MIDDLE CHILDHOOD						ADOLESCENCE													
	Prenatal ²	Newborn ³	3-5 d ⁴	By 1 mo	2 mo	4 mo	6 mo	9 mo	12 mo	15 mo	18 mo	24 mo	30 mo	3 y	4 y	5 y	6 y	7 y	8 y	9 y	10 y	11 y	12 y	13 y	14 y	15 y	16 y	17 y	18 y	19 y	20 y	21 y		
HISTORY Initial/Interval	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	
MEASUREMENTS																																		
Length/Height and Weight		●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	
Head Circumference		●	●	●	●	●	●	●	●	●	●	●	●	●	●																			
Weight for Length		●	●	●	●	●	●	●	●	●	●	●	●	●	●																			
Body Mass Index ⁵												●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	
Blood Pressure ⁶		★	★	★	★	★	★	★	★	★	★	★	★	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	
SENSORY SCREENING																																		
Vision ⁷		★	★	★	★	★	★	★	★	★	★	★	★	●	●	●	●	★	●	★	●	★	●	★	★	●	★	★	★	★	★	★	★	
Hearing		● ⁸	★	★	★	★	★	★	★	★	★	★	★	★	●	●	●	★	●	★	●	★	★	★	★	★	★	★	★	★	★	★	★	
DEVELOPMENTAL/BEHAVIORAL ASSESSMENT																																		
Developmental Screening ⁹								●				●	●																					
Autism Screening ¹⁰											●	●																						
Developmental Surveillance		●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	
Psychosocial/Behavioral Assessment		●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	
Alcohol and Drug Use Assessment ¹¹																						★	★	★	★	★	★	★	★	★	★	★	★	
Depression Screening ¹²																						●	●	●	●	●	●	●	●	●	●	●	●	
PHYSICAL EXAMINATION¹³		●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	
PROCEDURES¹⁴																																		
Newborn Blood Screening ¹⁵		←	●	→																														
Critical Congenital Heart Defect Screening ¹⁶		●																																
Immunization ¹⁷		●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	
Hematocrit or Hemoglobin ¹⁸					★				●	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	
Lead Screening ¹⁹						★	★	● or ★ ²⁰		★	● or ★ ²⁰		★	★	★	★	★																	
Tuberculosis Testing ²¹				★		★		★	★		★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	
Dyslipidemia Screening ²²												★		★		★	★		★	←	●	→	★	★	★	★	★	★	★	★	★	★	★	
STI/HIV Screening ²³																						★	★	★	★	★	★	★	★	★	★	★	★	★
Cervical Dysplasia Screening ²⁴																																	●	
ORAL HEALTH²⁵						★	★	● or ★		● or ★	● or ★	● or ★	●				●																	
Fluoride Varnish ²⁶							←	→	←	→	←	→	←	→	←	→																		
ANTICIPATORY GUIDANCE	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	

- If a child comes under care for the first time at any point on the schedule, or if any items are not accomplished at the suggested age, the schedule should be brought up to date at the earliest possible time.
- A prenatal visit is recommended for parents who are at high risk, for first-time parents, and for those who request a conference. The prenatal visit should include anticipatory guidance, pertinent medical history, and a discussion of benefits of breastfeeding and planned method of feeding, per the 2009 AAP statement "The Prenatal Visit" (<http://pediatrics.aappublications.org/content/124/4/1227.full>).
- Every infant should have a newborn evaluation after birth, and breastfeeding should be encouraged (and instruction and support should be offered).
- Every infant should have an evaluation within 3 to 5 days of birth and within 48 to 72 hours after discharge from the hospital to include evaluation for feeding and jaundice. Breastfeeding infants should receive formal breastfeeding evaluation, and their mothers should receive encouragement and instruction, as recommended in the 2012 AAP statement "Breastfeeding and the Use of Human Milk" (<http://pediatrics.aappublications.org/content/129/3/e827.full>). Newborn infants discharged less than 48 hours after delivery must be examined within 48 hours of discharge, per the 2010 AAP statement "Hospital Stay for Healthy Term Newborns" (<http://pediatrics.aappublications.org/content/125/2/405.full>).
- Screen, per the 2007 AAP statement "Expert Committee Recommendations Regarding the Prevention, Assessment, and Treatment of Child and Adolescent Overweight and Obesity: Summary Report" (http://pediatrics.aappublications.org/content/120/Supplement_4/S164.full).
- Blood pressure measurement in infants and children with specific risk conditions should be performed at visits before age 3 years.
- A visual acuity screen is recommended at ages 4 and 5 years, as well as in cooperative 3 year olds. Instrument based screening may be used to assess risk at ages 12 and 24 months, in addition to the well visits at 3 through 5 years of age. See 2016 AAP statement, "Visual System Assessment in Infants, Children, and Young Adults by Pediatricians" (<http://pediatrics.aappublications.org/content/137/1/1.51>) and "Procedures for Evaluation of the Visual System by Pediatricians" (<http://pediatrics.aappublications.org/content/137/1/1.52>).
- All newborns should be screened, per the AAP statement "Year 2007 Position Statement: Principles and Guidelines for Early Hearing Detection and Intervention Programs" (<http://pediatrics.aappublications.org/content/120/4/898.full>).
- See 2006 AAP statement "Identifying Infants and Young Children With Developmental Disorders in the Medical Home: An Algorithm for Developmental Surveillance and Screening" (<http://pediatrics.aappublications.org/content/118/1/405.full>).
- Screening should occur per the 2007 AAP statement "Identification and Evaluation of Children with Autism Spectrum Disorders" (<http://pediatrics.aappublications.org/content/120/5/1183.full>).

- A recommended screening tool is available at <http://www.ceasar-boston.org/CRAFFT/index.php>.
- Recommended screening using the Patient Health Questionnaire (PHQ)-2 or other tools available in the GLAD-PC toolkit and at http://www.aap.org/en-us/advocacy-and-policy/aap-health-initiatives/Mental-Health/Documents/MH_ScreeningChart.pdf.
- At each visit, age-appropriate physical examination is essential, with infant totally unclothed and older children undressed and suitably draped. See 2011 AAP statement "Use of Chaperones During the Physical Examination of the Pediatric Patient" (<http://pediatrics.aappublications.org/content/127/5/991.full>).
- These may be modified, depending on entry point into schedule and individual need.
- The Recommended Uniform Newborn Screening Panel (<http://www.hrsa.gov/advisorycommittees/mchbadvisory/heritableorders/recommendedpanel/uniformscreeningpanel.pdf>), as determined by The Secretary's Advisory Committee on Heritable Disorders in Newborns and Children, and state newborn screening laws/regulations (<http://genes-r-us.uhscsa.edu/sites/genes-r-us/files/nbsdisorders.pdf>), establish the criteria for and coverage of newborn screening procedures and programs. Follow-up must be provided, as appropriate, by the pediatrician.
- Screening for critical congenital heart disease using pulse oximetry should be performed in newborns, after 24 hours of age, before discharge from the hospital, per the 2011 AAP statement "Endorsement of Health and Human Services Recommendation for Pulse Oximetry Screening for Critical Congenital Heart Disease" (<http://pediatrics.aappublications.org/content/129/1/190.full>).
- Schedules, per the AAP Committee on Infectious Diseases, are available at: <http://aapredbook.aappublications.org/site/resources/zschedules.xhtml>. Every visit should be an opportunity to update and complete a child's immunizations.
- See 2010 AAP statement "Diagnosis and Prevention of Iron Deficiency and Iron Deficiency Anemia in Infants and Young Children (0-3 Years of Age)" (<http://pediatrics.aappublications.org/content/126/5/1040.full>).
- For children at risk of lead exposure, see the 2012 CDC Advisory Committee on Childhood Lead Poisoning Prevention statement "Low Level Lead Exposure Harms Children: A Renewed Call for Primary Prevention" (http://www.cdc.gov/nceh/lead/ACCLPP/Final_Document_030712.pdf).
- Perform risk assessments or screenings as appropriate, based on universal screening requirements for patients with Medicaid or in high prevalence areas.

- Tuberculosis testing per recommendations of the Committee on Infectious Diseases, published in the current edition of *AAP Red Book: Report of the Committee on Infectious Diseases*. Testing should be performed on recognition of high-risk factors.
- See AAP-endorsed 2011 guidelines from the National Heart Blood and Lung Institute, "Integrated Guidelines for Cardiovascular Health and Risk Reduction in Children and Adolescents" (http://www.nhlbi.nih.gov/guidelines/cvd_ped/index.htm).
- Adolescents should be screened for sexually transmitted infections (STIs) per recommendations in the current edition of the *AAP Red Book: Report of the Committee on Infectious Diseases*. Additionally, all adolescents should be screened for HIV according to the AAP statement (<http://pediatrics.aappublications.org/content/128/5/1023.full>) once between the ages of 16 and 18, making every effort to preserve confidentiality of the adolescent. Those at increased risk of HIV infection, including those who are sexually active, participate in injection drug use, or are being tested for other STIs, should be tested for HIV and reassessed annually.
- See USPSTF recommendations (<http://www.uspreventiveservicestaskforce.org/uspstf/uspstf.htm>). Indications for pelvic examinations prior to age 21 are noted in the 2010 AAP statement "Gynecologic Examination for Adolescents in the Pediatric Office Setting" (<http://pediatrics.aappublications.org/content/126/3/583.full>).
- Assess if the child has a dental home. If no dental home is identified, perform a risk assessment (<http://www2.aap.org/oralhealth/docs/RiskAssessmentTool.pdf>) and refer to a dental home. If primary water source is deficient in fluoride, consider oral fluoride supplementation. Recommend brushing with fluoride toothpaste in the proper dosage for age. See 2009 AAP statement "Oral Health Risk Assessment Timing and Establishment of the Dental Home" (<http://pediatrics.aappublications.org/content/111/5/1113.full>), 2014 clinical report "Fluoride Use in Caries Prevention in the Primary Care Setting" (<http://pediatrics.aappublications.org/content/134/3/626>), and 2014 AAP statement "Maintaining and Improving the Oral Health of Young Children" (<http://pediatrics.aappublications.org/content/134/6/1224.full>).
- See USPSTF recommendations (<http://www.uspreventiveservicestaskforce.org/uspstf/uspstf.htm>). Once teeth are present, fluoride varnish may be applied to all children every 3-6 months in the primary care or dental office. Indications for fluoride use are noted in the 2014 AAP clinical report "Fluoride Use in Caries Prevention in the Primary Care Setting" (<http://pediatrics.aappublications.org/content/134/3/626>).

**Summary of changes made to the
Bright Futures/AAP Recommendations for Preventive Pediatric Health Care**
(Periodicity Schedule)

This Schedule reflects changes approved in October 2015 and published in January 2016. For updates, visit www.aap.org/periodicityschedule.

Changes made October 2015

- **Vision Screening-** The routine screening at age 18 has been changed to a risk assessment.
- Footnote 7 has been updated to read, “A visual acuity screen is recommended at ages 4 and 5 years, as well as in cooperative 3 year olds. Instrument based screening may be used to assess risk at ages 12 and 24 months, in addition to the well visits at 3 through 5 years of age. See 2016 AAP statement, “Visual System Assessment in Infants, Children, and Young Adults by Pediatricians (<http://pediatrics.aappublications.org/content/137/1/1.51>) and “Procedures for Evaluation of the Visual System by Pediatricians” (<http://pediatrics.aappublications.org/content/137/1/1.52>).

Changes made May 2015

- **Oral Health-** A subheading has been added for fluoride varnish, with a recommendation from 6 months through 5 years.
- Footnote 25 wording has been edited and also includes reference to the 2014 clinical report, “Fluoride Use in Caries Prevention in the Primary Care Setting” (<http://pediatrics.aappublications.org/content/134/3/626>) and 2014 policy statement, “Maintaining and Improving the Oral Health of Young Children” (<http://pediatrics.aappublications.org/content/134/6/1224.full>).
- Footnote 26 has been added to the new fluoride varnish subheading: See USPSTF recommendations (<http://www.uspreventiveservicestaskforce.org/uspstf/uspsdnch.htm>). Once teeth are present, fluoride varnish may be applied to all children every 3-6 months in the primary care or dental office. Indications for fluoride use are noted in the 2014 AAP clinical report “Fluoride Use in Caries Prevention in the Primary Care Setting” (<http://pediatrics.aappublications.org/content/134/3/626>).

Changes made March 2014

Changes to Developmental/Behavioral Assessment

- **Alcohol and Drug Use Assessment-** Information regarding a recommended screening tool (CRAFFT) was added.
- **Depression-** Screening for depression at ages 11 through 21 has been added, along with suggested screening tools.

Changes to Procedures

- **Dyslipidemia screening-** An additional screening between 9 and 11 years of age has been added. The reference has been updated to the AAP-endorsed National Heart Blood and Lung Institute policy (http://www.nhlbi.nih.gov/guidelines/cvd_ped/index.htm).
- **Hematocrit or hemoglobin-** A risk assessment has been added at 15 and 30 months. The reference has been updated to the current AAP policy (<http://pediatrics.aappublications.org/content/126/5/1040.full>).
- **STI/HIV screening-** A screen for HIV has been added between 16 and 18 years. Information on screening adolescents for HIV has been added in the footnotes. STI screening now references recommendations made in the AAP Red Book. This category was previously titled “STI Screening.”
- **Cervical dysplasia-** Adolescents should no longer be routinely screened for cervical dysplasia until age 21. Indications for pelvic exams prior to age 21 are noted in the 2010 AAP statement “Gynecologic Examination for Adolescents in the Pediatric Office Setting” (<http://pediatrics.aappublications.org/content/126/3/583.full>).
- **Critical Congenital Heart Disease-** Screening for critical congenital heart disease using pulse oximetry should be performed in newborns, after 24 hours of age, before discharge from the hospital, per the 2011 AAP statement, “Endorsement of Health and Human Services Recommendation for Pulse Oximetry Screening for Critical Congenital Heart Disease” (<http://pediatrics.aappublications.org/content/129/1/190.full>).

See www.aap.org/periodicityschedule for additional updates made to footnotes and references in March 2014.