

WHERE IS THE VIS DATE LOCATED?

ANSWER: The "VIS Date" is located at the bottom of the first OR second page of the Vaccine Information Statement (VIS) depending on the type of vaccine.

VACCINE INFORMATION STATEMENT

Tdap Vaccine

What You Need to Know

(Tetanus, Diphtheria and Pertussis)

Many Vaccine Information Statements are available in Spanish and other languages. See www.immunize.org/vis
Hojas de información sobre vacunas están disponibles en español y en muchos otros idiomas. Visite www.immunize.org/vis

1 Why get vaccinated?

Tetanus, diphtheria and pertussis are very serious diseases. Tdap vaccine can protect us from these diseases. And, Tdap vaccine given to pregnant women can protect newborn babies against pertussis.

TETANUS (Lockjaw) is rare in the United States today. It causes painful muscle tightening and stiffness, usually all over the body.

- It can lead to tightening of muscles in the head and neck so you can't open your mouth, swallow, or sometimes even breathe. Tetanus kills about 1 out of 10 people who are infected even after receiving the best medical care.

DIPHTHERIA is also rare in the United States today. It can cause a thick coating to form in the back of the throat.

2 Tdap vaccine

Tdap vaccine can protect adolescents and adults from tetanus, diphtheria, and pertussis. One dose of Tdap is routinely given at age 11 or 12. People who did *not* get Tdap at that age should get it as soon as possible.

Tdap is especially important for healthcare professionals and anyone having close contact with a baby younger than 12 months.

Pregnant women should get a dose of Tdap during **every pregnancy**, to protect the newborn from pertussis. Infants are most at risk for severe, life-threatening complications from pertussis.

Another vaccine, called Td, protects against tetanus and diphtheria, but not pertussis. A Td booster should be given every 10 years. Tdap may be given as one of these boosters if you have never gotten Tdap before. Tdap may also be given after a severe cut or burn to prevent

(First Page)

5 people in 100)

- Swelling of the entire arm where the shot was given (up to about 1 in 500).

Severe problems following Tdap
(Unable to perform usual activities; required medical attention)

- Swelling, severe pain, bleeding and redness in the arm where the shot was given (rare).

Problems that could happen after any vaccine:

- People sometimes faint after a medical procedure, including vaccination. Sitting or lying down for about 15 minutes can help prevent fainting, and injuries caused by a fall. Tell your doctor if you feel dizzy, or have vision changes or ringing in the ears.
- Some people get severe pain in the shoulder and have difficulty moving the arm where a shot was given. This happens very rarely.
- Any medication can cause a severe allergic reaction. Such reactions from a vaccine are very rare, estimated at fewer than 1 in a million doses, and would happen within a few minutes to a few hours after the vaccination.

Persons who believe they may have been injured by a vaccine can learn about the program and about filing a claim by calling 1-800-338-2382 or visiting the VICP website at www.hrsa.gov/vaccinecompensation. There is a time limit to file a claim for compensation.

7 How can I learn more?


- Ask your doctor. He or she can give you the vaccine package insert or suggest other sources of information.
- Call your local or state health department.
- Contact the Centers for Disease Control and Prevention (CDC):
 - Call 1-800-232-4636 (1-800-CDC-INFO) or
 - Visit CDC's website at www.cdc.gov/vaccines

Vaccine Information Statement
Tdap Vaccine

8/6/21

42 U.S.C. § 300aa-26

Office Use Only



(Second Page)