

Subject: Platelet-rich Plasma (PRP)		Original Effective Date: 10/8/2014					
Policy Number:	Revision Date(s):	10/8/2014					
MCP-207	Revision Date(s).						
	0/12/2019 6/19/2019 6/17/2020	4/5/01					
Review Date: 12/16/2015, 9/15/2016, 9/19/2017,	, , , , , , , , , , , , , , , , , , , ,	4/5/21					
MCPC Approval Date: 9/19/2017, 9/13/2018, 6/2	19/2019, 6/17/2020, 4/5/21	MCPC Approval Date: 9/19/2017, 9/13/2018, 6/19/2019, 6/17/2020, 4/5/21					

Contents

DISCLAIMER	1
DESCRIPTION OF PROCEDURE/SERVICE/PHARMACEUTICAL	1
Position Statement	
Summary of Medical Evidence	
Coding Information	
References	
CENTERS FOR MEDICARE AND MEDICAID SERVICES (CMS)	
Review/Revision History	

DISCLAIMER

This Molina Clinical Policy (MCP) is intended to facilitate the Utilization Management process. It expresses Molina's determination as to whether certain services or supplies are medically necessary, experimental, investigational, or cosmetic for purposes of determining appropriateness of payment. The conclusion that a particular service or supply is medically necessary does not constitute a representation or warranty that this service or supply is covered (i.e., will be paid for by Molina) for a particular member. The member's benefit plan determines coverage. Each benefit plan defines which services are covered, which are excluded, and which are subject to dollar caps or other limits. Members and their providers will need to consult the member's benefit plan to determine if there are any exclusion(s) or other benefit limitations applicable to this service or supply. If there is a discrepancy between this policy and a member's plan of benefits, the benefits plan will govern. In addition, coverage may be mandated by applicable legal requirements of a State, the Federal government or CMS for Medicare and Medicaid members. CMS's Coverage Database can be found on the CMS website. The coverage directive(s) and criteria from an existing National Coverage Determination (NCD) or Local Coverage Determination (LCD) will supersede the contents of this Molina Clinical Policy (MCP) document and provide the directive for all Medicare members.

DESCRIPTION OF PROCEDURE/SERVICE/PHARMACEUTICAL

Platelet-rich plasma (PRP) is a blood product derived from plasma that contains an increased concentration of platelets. PRP is also referred to as autologous platelet concentrate (APC) and autologous platelet gel (APG). The use of PRP is an approach being investigated for the treatment of soft tissue and bone healing, chronic non-healing wounds including burns and diabetic ulcers, osteoarthritis, tendon and ligament injuries and other surgeries. It is proposed that activated platelets initiate repair by releasing potent locally acting growth factors



that stimulate a connective tissue response, causing division and migration of fibroblasts and formation of new capillaries to aid in the healing process. Platelet-rich plasma is usually prepared by a clinician or technician where blood is taken from the patient and centrifuged to obtain a concentrated suspension of platelets. PRP is injected or implanted during surgery with the goal of accelerating healing of the damaged tendon or ligament. For wound healing PRP is applied directly to the wound surface to promote growth of skin, soft tissue, and blood vessels.

POSITION STATEMENT 2-	-25	2-	4	г	'n	V	יןי	G	n	./	۸	7	L	r	r	٨	r	ļ	C		N	N	c	T	7	1	C	n)	1
-----------------------	-----	----	---	---	----	---	-----	---	---	----	---	---	---	---	---	---	---	---	---	--	---	---	---	---	---	---	---	---	---	---

	t rich plasma is considered experimental, investigational and unproven because of insufficient ce in the peer reviewed medical literature for any of the following conditions:
	Achilles tendon repair
	Acute surgical wounds
	Anterior cruciate ligament repair
	Burns
	Chronic non-healing diabetic, venous and/or pressure wounds*
	Dupuytren's contracture
	Epicondylitis (e.g., tennis elbow, elbow epicondylar tendinosis)
	Hamstring tendon injuries
	Injection of any ligament or tendon injury
	Osteoarthritis of the hip & knee
	Periodontal surgery
	Plantar fasciitis
	Rotator cuff repair
	Sinus surgery
	Spinal fusion or any other surgery using bone grafting
	Various tendinopathies
	All other conditions not listed above

*Note: See CMS NCD for Medicare members 1

SUMMARY OF MEDICAL EVIDENCE 3-34

Results from both randomized controlled trials (RCTs) and nonrandomized controlled studies provide varied and inconclusive evidence regarding the ability of injection of platelet-rich plasma (PRP) to improve outcomes or accelerate healing in patients who have tendon or ligament injuries. There is insufficient published evidence to assess the safety and/or impact on health outcomes or patient management for platelet rich plasma for any indication. Below is a summary of the most relevant evidence based studies.

Chronic Wounds

According to the Cochrane Review (2012) there is currently no evidence to suggest that autologous PRP is of value for treating chronic wounds. The reports analyzed were based on small numbers of randomized controlled



studies for the treatment of chronic wounds including 325 patients, most of whom were at either high or unclear risk of bias. ³

A systematic review and meta-analysis evaluated the use platelet rich plasma (PRP) for the treatment of cutaneous wounds compared to standard wound care. These studies included 3 systematic reviews, 12 randomized controlled trials, 2 prospective cohort studies, 3 prospective comparative studies and 4 retrospective reviews. The results of the meta-analysis suggested that PRP therapy can positively impact wound healing and associated factors such as pain and infection in cutaneous wounds. Limitations of the studies included heterogeneous patient populations, lack of long-term follow-up, and pooling of data on different types of PFG products and regimens. Several of the studies included in the meta-analysis had conflicting results. ⁴

Knee Osteoarthritis

A meta-analysis in a systematic review of 6 studies, including 577 patients, compared the outcomes of patients with symptomatic knee osteoarthritis treated by platelet-rich plasma, hyaluronic acid or normal saline (placebo). There was no difference in the pooled results for visual analog scale score or overall patient satisfaction.

Adverse events occurred more frequently in patients treated with PRP than in those treated with HA/placebo. ⁵ In a RCT of 109 patients with knee degenerative pathology treated by platelet-rich plasma (n=54) or hyaluronic acid injections (n=55), there was no statistically significant differences observed between groups at 12-month follow-up. ⁶ Another RCT of 78 patients with bilateral OA were divided randomly into 3 groups. Group A (52 knees) received a single injection of PRP, group B (50 knees) received 2 injections of PRP 3 weeks apart, and group C (46 knees) received a single injection of normal saline. Results reported that a single dose of WBC-filtered PRP in concentrations of 10 times the normal amount is as effective as 2 injections to alleviate symptoms in early knee OA. The results, however, deteriorate after 6 months. ²¹ RCT's compared the effectiveness of intraarticular (IA) multiple and single platelet-rich plasma (PRP) injections as well as hyaluronic acid (HA) injections in different stages of osteoarthritis (OA) of the knee and found there was no significant difference in the scores of patients injected with one dose of PRP or HA. ²⁴⁻²⁵

Tendon and Ligament Injuries

A long-term pilot study intratendinous injection of platelet-rich plasma under US guidance to treat tendinopathy in the upper (medial and lateral epicondylar tendons) and the lower (patellar, Achilles, hamstring and adductor longus, and peroneal tendons) limbs of 408 patients reported that residual US size of lesions were lower after intratendinous injection of PRP under US guidance at 6 weeks and during long-term follow-up (32 weeks) compared with baseline. ⁷

There are several RCTs that evaluated PRP for tendon and ligament injuries. All studies found PRP treatment to be reasonably safe. Although many of the RCTs reported double- or single-blinding, all of the studies were relatively small, with treatment and control groups that had 10 to 80 patients and evaluated PRP as an adjunct to surgery for treatment of anterior cruciate ligament (ACL) injuries ⁸⁻¹⁰ Other RCTs evaluated PRP as an adjunct to arthroscopic or open surgery for the treatment of rotator cuff injuries ¹¹⁻¹³ ²² and chronic rotator cuff tendinopathy. ²⁰ Three RCTs evaluated PRP for the treatment of elbow tendon injuries, such as lateral epicondylitis or elbow tendinopathy. ¹⁴⁻¹⁶ Several RCTs evaluated PRP for the treatment of Achilles tendinopathy or tendon rupture. ¹⁷ ¹⁸ ²³ One RCT assessed PRP in hamstring injuries. ¹⁹ Results from these RCTs



provide mixed and inconclusive evidence regarding the ability of injection of platelet-rich plasma (PRP) to improve outcomes or accelerate healing in patients who have tendon or ligament injuries. A Cochrane review (2014) found there is currently insufficient evidence to support the use of PRT for treating musculoskeletal soft tissue injuries. ³

CODING INFORMATION: THE CODES LISTED IN THIS POLICY ARE FOR REFERENCE PURPOSES ONLY. LISTING OF A SERVICE OR DEVICE CODE IN THIS POLICY DOES NOT IMPLY THAT THE SERVICE DESCRIBED BY THIS CODE IS A COVERED OR NON-COVERED. COVERAGE IS DETERMINED BY THE BENEFIT DOCUMENT. THIS LIST OF CODES MAY NOT BE ALL INCLUSIVE.

CPT	Description
0232T	Injection(s), platelet rich plasma, any tissue, including image guidance, harvesting and
	preparation when performed

HCPCS	Description
G0460	Autologous platelet rich plasma for chronic wounds/ulcers, including phlebotomy, centrifugation,
	and all other preparatory procedures, administration and dressings, per treatment
P9020	Platelet rich plasma, each unit

ICD-1	Description: [For dates of service on or after 10/01/2015]
	Any Diagnosis

REFERENCES

Government Agency

1. Centers for Medicare & Medicaid Services (CMS). Medicare Coverage Database. National coverage determination (NCD) for Blood-Derived Products for Chronic Non-Healing Wounds (270.3). Effective 8/2/2012. Accessed at: http://www.cms.gov/medicare-coverage-database/search/advanced-search.aspx.

Hayes

- 2. Hayes a TractManager Company. Winifred Hayes Inc., Lansdale, PA
 - Comparative Effectiveness Review: Platelet-Rich Plasma for Treatment of Conditions of the Achilles Tendon and Plantar Fascia. March 2018, updated June, 2020.
 - Comparative Effectiveness Review: Platelet-Rich Plasma for Rotator Cuff Repairs, Tendinopathies, and Related Conditions: A Review of Reviews. June 2019, Updated Sep, 2020.
 - Comparative Effectiveness Review: Platelet-Rich Plasma for Treatment of Lateral Epicondylitis: A Review of Reviews. Dec, 2017. Updated Dec, 2019.
 - Comparative Effectiveness Review: Platelet-Rich Plasma for Knee Osteoarthritis: A Review of Reviews. Nov 2017, updated Dec 2019.
 - Comparative Effectiveness Review: Platelet-Rich Plasma for Tendinopathies Or Ligament Injuries Of The Knee. Dec 2017, updated Dec 2020.
 - Platelet-Rich Plasma for Treatment of Thoracic and Lumbar Spinal Pain. Oct 2018. [Archived Nov 2019]
 - HTA: Platelet-Rich Plasma for Hip Osteoarthritis. June, 2019. Updated Oct, 2020



Peer Reviewed Literature

3. Cochrane:

- Martinez-Zapata MJ, Marti-Carvajal A, Solà I, Autologous platelet-rich plasma for treating chronic wounds. Cochrane Database Syst Rev, 17. 2012. CD006899.
- Moraes VY, Lenza M, et al. Platelet-rich therapies for musculoskeletal soft tissue injuries.
 Cochrane Database Syst Rev. 2014 Apr 29;(4):CD010071. doi: 10.1002/14651858.CD010071.pub3.
- 4. Carter MJ, Fylling CP, Parnell LK. Use of platelet rich plasma gel on wound healing: a systematic review and meta-analysis. Eplasty. 2011;11:e38.
- 5. Khoshbin A, Leroux T, Wasserstein D, Marks P, Theodoropoulos J et al. The efficacy of platelet-rich plasma in the treatment of symptomatic knee osteoarthritis: a systematic review with quantitative synthesis. Arthroscopy. 2013;29(12): 2037-48
- 6. Filardo G, Kon E, Di Martino A, Di Mattio B et al. Platelet-rich plasma vs hyaluronic acid to treat knee degenerative pathology: study design and preliminary results of a randomized controlled trial. BMC musculoskeletal disorders. 2012;13: 229
- 7. Dallaudiere B., Pesquer L., Meyer P et al. Intratendinous injection of platelet-rich plasma under US guidance to treat tendinopathy: A long-term pilot study. Journal of Vascular and Interventional Radiology. 25 (5) (pp 717-723), 2014.
- 8. Cervellin M, de GL, Bait C, Denti M, Volpi P. Autologous platelet-rich plasma gel to reduce donor-site morbidity after patellar tendon graft harvesting for anterior cruciate ligament reconstruction: a randomized, controlled clinical study. Knee Surg Sports Traumatol Arthrosc. 2012;20(1):114-120
- 9. de Almeida AM, Demange MK, Sobrado MF et al. Patellar tendon healing with platelet-rich plasma: a prospective randomized controlled trial. Am J Sports Med. 2012;40(6):1282-1288.
- 10. Vogrin M, Rupreht M, Crnjac A et al. The effect of platelet-derived growth factors on knee stability after anterior cruciate ligament reconstruction: a prospective randomized clinical study. Wien Klin Wochenschr. 2010b;122(Suppl 2):91-95.
- 11. Rodeo SA, Delos D, Williams RJ et al. The effect of platelet-rich fibrin matrix on rotator cuff tendon healing: a prospective, randomized clinical study. Am J Sports Med. 2012;40(6):1234-1241.
- 12. Randelli P, Arrigoni P, Ragone V et al. Platelet rich plasma in arthroscopic rotator cuff repair: a prospective RCT study, 2-year follow-up. J Shoulder Elbow Surg. 2011;20(4):518-528.
- 13. Gumina S, Campagna V, Ferrazza G, et al. Use of platelet-leukocyte membrane in arthroscopic repair of large rotator cuff tears: a prospective randomized study. J Bone Joint Surg Am. 2012;94(15):1345-1352.
- 14. Peerbooms JC, Sluimer J, Bruijn DJ, Gosens T. Positive effect of an autologous platelet concentrate in lateral epicondylitis in a double-blind randomized controlled trial: platelet-rich plasma versus corticosteroid injection with a 1-year follow-up. Am J Sports Med. 2010;38(2):255-262.
- 15. Creaney L, Wallace A, Curtis M, Connell D. Growth factor-based therapies provide additional benefit beyond physical therapy in resistant elbow tendinopathy: a prospective, single-blind, randomised trial of autologous blood injections versus platelet-rich plasma injections. Br J Sports Med. 2011;45(12):966-971.
- 16. Thanasas C, Papadimitriou G, Charalambidis C et al. Platelet-rich plasma versus autologous whole blood for the treatment of chronic lateral elbow epicondylitis: a randomized controlled clinical trial. Am J Sports Med. 2011;39(10):2130-2134.



- 17. de Vos RJ, Weir A, van Schie HT, et al. Platelet-rich plasma injection for chronic Achilles tendinopathy: a randomized controlled trial. JAMA. 2010;303(2):144-149.
- 18. de Jonge S, de Vos RJ, Weir A, et al. One-year follow-up of platelet-rich plasma treatment in chronic Achilles tendinopathy: a double-blind randomized placebo-controlled trial. Am J Sports Med. 2011;39(6):1623-1629.
- 19. Reurink G1, Goudswaard GJ, Moen MH et al. Platelet-rich plasma injections in acute muscle injury. N Engl J Med. 2014 Jun 26;370(26):2546-7. doi: 10.1056/NEJMc1402340.
- 20. Kesikburun S1, Tan AK et al. Platelet-rich plasma injections in the treatment of chronic rotator cuff tendinopathy: a randomized controlled trial with 1-year follow-up. Am J Sports Med. 2013 Nov;41(11):2609-16. doi: 10.1177/0363546513496542. Epub 2013 Jul 26.
- 21. Patel S1, Dhillon MS et al. Treatment with platelet-rich plasma is more effective than placebo for knee osteoarthritis: a prospective, double-blind, randomized trial. Am J Sports Med. 2013 Feb;41(2):356-64. doi: 10.1177/0363546512471299. Epub 2013 Jan 8.
- 22. Flury M, Rickenbacher D, Schwyzer HK et al. Does Pure Platelet-Rich Plasma Affect Postoperative Clinical Outcomes After Arthroscopic Rotator Cuff Repair? A Randomized Controlled Trial. Am J Sports Med. 2016 Aug;44(8):2136-46. doi: 10.1177/0363546516645518. Epub 2016 May 16.
- 23. Krogh TP, Ellingsen T, Christensen R et al. Ultrasound-Guided Injection Therapy of Achilles Tendinopathy With Platelet-Rich Plasma or Saline: A Randomized, Blinded, Placebo-Controlled Trial. Am J Sports Med. 2016 Aug;44(8):1990-7. doi: 10.1177/0363546516647958. Epub 2016 Jun 2.
- 24. Görmeli G, Görmeli CA, et al. Multiple PRP injections are more effective than single injections and hyaluronic acid in knees with early osteoarthritis: a randomized, double-blind, placebo-controlled trial. Knee Surg Sports Traumatol Arthrosc. 2017 Mar;25(3):958-965. doi: 10.1007/s00167-015-3705-6. Epub 2015 Aug 2.
- 25. Montañez-Heredia E, Irízar S, et al. Intra-articular injections of platelet-rich plasma versus hyaluronic acid in the treatment of osteoarthritic knee pain: a randomized clinical trial in the context of the Spanish National Health Care System. Int J Mol Sci. 2016 Jul 2;17(7). pii: E1064. doi: 10.3390/ijms17071064.
- 26. Kim S, Hwang J, Kim MJ, Lim JY, Lee WH, Choi JE. Systematic review with network meta-analysis of randomized controlled trials of rotator cuff tear treatment. Int J Technol Assess Health Care. 2018a:34(1):78-86.
- 27. Kim SJ, Kim EK, Kim SJ, Song DH. Effects of bone marrow aspirate concentrate and platelet-rich plasma on patients with partial tear of the rotator cuff tendon. J Orthop Surg Res. 2018b;13(1):1.
- 28. Hurley ET, Lim Fat D, Moran CJ, Mullett H. The efficacy of platelet-rich plasma and platelet-rich fibrin in arthroscopic rotator cuff repair: a meta-analysis of randomized controlled trials. Am J Sports Med. 2018:363546517751397. Epub ahead of print. February 21, 2018. Available at: http://journals.sagepub.com/doi/pdf/10.1177/0363546517751397.
- 29. Andia I, Maffulli N. Update on platelet-rich plasma for shoulder and elbow tendinopathy. Tech Shoulder Elb Surg. 2017;18(3):91-100.
- 30. Filardo G, Di Matteo B, Kon E, Merli G, Marcacci M. Platelet-rich plasma in tendon-related disorders: results and indications. Knee Surg Sports Traumatol Arthrosc. 2016. Epub ahead of print. September 24, 2016. Available at: https://link.springer.com/article/10.1007%2Fs00167-016-4261-4. Accessed November 7, 2017.



- 31. Fitzpatrick J, Bulsara MK, McCrory PR, Richardson MD, Zheng MH. Analysis of platelet-rich plasma extraction: variations in platelet and blood components between 4 common commercial kits. Orthop J Sports Med. 2017a;5(1):2325967116675272.
- 32. Fitzpatrick J, Bulsara M, Zheng MH. The effectiveness of platelet-rich plasma in the treatment of tendinopathy: a meta-analysis of randomized controlled clinical trials. Am J Sports Med. 2017b;45(1):226-233.
- 33. Mi B, Liu G, Zhou W, et al. Platelet rich plasma versus steroid on lateral epicondylitis: meta-analysis of randomized clinical trials. Phys Sportsmed. 2017;45(2):97-104.
- 34. Schoffl V, Willauschus W, Sauer F, et al. Autologous conditioned plasma versus placebo injection therapy in lateral epicondylitis of the elbow: a double blind, randomized study. Sportverletz Sportschaden. 2017;31(1):31-36.

Professional Society Guidelines

- 35. National Institute for Healthcare and Excellence (NICE). Platelet-rich plasma injections for osteoarthritis of the knee. Interventional procedures guidance [IPG491] Published date: May 2014
- 36. American Academy of Orthopedic Surgeons (AAOS). Accessed at: https://www.aaos.org/
 - Appropriate use criteria for non-arthroplasty treatment of osteoarthritis of the knee. 2015. [archived]
 - Clinical practice guideline on the treatment of osteoarthritis of the knee, 2nd edition. 2015. [archived]
 - OrthoInfo. Platelet-Rich Plasma (PRP). 2011. Accessed at: http://orthoinfo.aaos.org/topic.cfm?topic=A00648

CENTERS FOR MEDICARE AND MEDICAID SERVICES (CMS)

The coverage directive(s) and criteria from an existing National Coverage Determination (NCD) or Local Coverage Determination (LCD) will supersede the contents of this Molina Clinical Policy (MCP) document and provide the directive for all Medicare members.

(CMS) has determined that platelet-rich plasma (PRP) an autologous blood-derived product, will be covered only for the treatment of chronic non-healing diabetic, venous and/or pressure wounds **only** when the patient is enrolled in a randomized clinical trial. ¹

REVIEW/REVISION HISTORY

10/8/14: New Policy

12/16/15, 9/15/16, 9/19/17: Policy reviewed, no changes

9/13/18, 6/19/19, 6/17/20: Policy reviewed, current literature indicates that procedure remains experimental, investigational and unproven for all indications. References updated.

4/5/21: Policy reviewed, no changes based on an evaluation of the literature that identified no new evidence to change our position.

