

DISCLAIMER

This Molina Clinical Policy (MCP) is intended to facilitate the Utilization Management process. Policies are not a supplementation or recommendation for treatment; Providers are solely responsible for the diagnosis, treatment, and clinical recommendations for the Member. It expresses Molina's determination as to whether certain services or supplies are medically necessary, experimental, investigational, or cosmetic for purposes of determining appropriateness of payment. The conclusion that a particular service or supply is medically necessary does not constitute a representation or warranty that this service or supply is covered (e.g., will be paid for by Molina) for a particular Member. The Member's benefit plan determines coverage – each benefit plan defines which services are covered, which are excluded, and which are subject to dollar caps or other limits. Members and their Providers will need to consult the Member's benefit plan to determine if there are any exclusion(s) or other benefit limitations applicable to this service or supply. If there is a discrepancy between this policy and a Member's plan of benefits, the benefits plan will govern. In addition, coverage may be mandated by applicable legal requirements of a State, the Federal government or CMS for Medicare and Medicaid Members. CMS's Coverage Database can be found on the CMS website. The coverage directive(s) and criteria from an existing National Coverage Determination (NCD) or Local Coverage Determination (LCD) will supersede the contents of this MCP and provide the directive for all Medicare members. References included were accurate at the time of policy approval and publication.

OVERVIEW

Hallux valgus (HV), also referred to as a bunion, is the most common foot deformity, with an estimated prevalence of 23% in people aged 18 to 65, and 35.7% in individuals over 65 (Ray et al. 2019). It is a deformity of the forefoot affecting the first ray and characterized by a medial bony protrusion of the first metatarsal head that frequently results in altered joint mechanics, functional impairment, and progressive foot discomfort (Dynamed 2025). The precise etiology is unknown but is thought to be multifaceted, involving genetics, joint hypermobility, aberrant foot anatomy/biomechanics, and footwear. Radiographs are recommended when surgery is being considered as a treatment option for severe or refractory symptoms to assess the severity of the deformity and determine the appropriate surgical approach. Primary treatment for HV should be conservative treatment, with most patients achieving sufficient symptomatic relief with nonsurgical interventions such as footwear modifications, orthoses, splinting, stretching, activity modifications, and/or manipulation, as well as the use of ice, analgesics, and bunion pads (Ettinger et al. 2025). Surgical management is reserved for patients who have not responded to nonsurgical treatment with the goals of pain relief, correction of the deformity, improved first-ray stability, and improved quality of life (Ferrari & Monson 2025).

There are several surgical options for treating HV, with the primary goal of realigning and balancing the first metatarsophalangeal (MTP) joint to relieve pain and improve quality of life. Generally, surgical treatment techniques are determined by the patient's condition, degree of deformity, symptoms, and radiological data. Radiological images visually quantify deformity severity using angles, such as the hallux valgus angle (HVA), the intermetatarsal angle (IMA), and, for certain patients, the distal metatarsal articular angle measurements.

HV may be sufficiently treated with exostectomy (e.g., surgical excision of bony growths), whereas more severe presentations may require osteotomy (e.g., bone repositioning) or arthrodesis (e.g., fusing bones together). Surgical operative management typically involves a combination of osseous and soft tissue procedures. Common bunion surgical procedures include:

- Exostectomy (bunionectomy), is the surgical excision of bony growths. It may be sufficient for treating simple bunions; however, it is usually performed with osteotomy and soft tissue repair since removing the bunion does not realign the joint. Procedures other than exostectomy should be considered for bunion treatment if the HVA is greater than 40 degrees on imaging.
- Osteotomy involves cutting the first metatarsal and realigning the bone in a less adducted position. The procedure repositions it to transfer load or weight-bearing from the pathologic to the normal joint alignment and is usually accompanied by soft tissue repair.
- Arthrodesis (e.g., bones fusion) is reserved for patients with severe arthritis or bunions, or those whose previous bunion surgery failed to solve the issue.

Bunionette deformity, also known as Taylor's bunion, involves the fifth metatarsal head with a painful lateral bony prominence. It is often associated with constrictive footwear causing pain, inflammation, keratosis, and ulceration. The fourth to fifth intermetatarsal angle varies between 3 and 11 degrees, with a mean of 6.5 to 8.0 degrees. The angle is often more than 10 degrees in patients with symptomatic bunionette deformity. The fifth metatarsophalangeal angle is

Molina Clinical Policy
Foot Surgery: Bunionectomy
Policy No. 700

Last Approval: 02/11/2026

Next Review Due By: February 2027



normally less than 14 degrees and is more than 16 degrees in those with bunionette (Shi et al. 2018). Nonsurgical therapy of symptomatic bunionette is frequently effective. However, when conservative management fails, surgical methods include condylar excision, and proximal or distal osteotomies.

RELATED POLICIES

MCP-701: Foot Surgery: Hallux Rigidus

MCP-702: Foot Surgery: Lesser Toe Deformities (Hammer, Mallet, and Claw Toe)

COVERAGE POLICY

Surgical correction of a bunion or bunionette deformity may be **considered medically necessary** when ALL the following criteria are met with accompanying documentation:

1. Member is \geq 18 years old, or has documented evidence of skeletal maturity
2. At least ONE of the following signs/symptoms directly attributable to a hallux valgus (HV) or bunionette deformity:
 - a. Significant and persistent pain at first metatarsophalangeal (MTP) joint or fifth MTP joint
 - b. Ulceration or skin breakdown at first MTP joint or fifth MTP joint
 - c. Clinically significant functional limitation resulting in impaired ambulation
3. Clinically significant symptoms resulting in persistent pain and functional limitation despite at least 6 months of provider directed conservative treatment (e.g., alternative or modified footwear, protective cushions or taping, foot orthotics, pharmacotherapy, and/or debridement or trimming of hyperkeratotic lesions)
4. Documentation of adequate lower extremity vascular perfusion (e.g., strong, palpable pedal pulses)
5. Diagnosis of HV or bunionette deformity with radiographic confirmation and interpretation via weight-bearing anterior/posterior and lateral views of the affected foot demonstrating ONE of the following:
 - a. For **Simple Bunionectomy** (i.e., removal of soft tissue and bony prominence without bony alignment correction):
 - i. Greater than 15 degrees hallux valgus angle (HVA) in the anteroposterior and lateral views
 - ii. No degenerative changes, absent or mild arthritis, at the MTP joint
 - b. For **Bunionectomy with Corrective Osteotomy**:
 - i. 20 degrees or greater HVA, indicating moderate to severe HV deformity
 - ii. 14 degrees or greater intermetatarsal angle (IMA), indicating moderate to severe deformity
 - iii. No degenerative changes, absent or mild arthritis, at the MTP joint
 - c. For **Bunionette Deformity** (i.e., Tailor's bunion) (e.g., osteotomy or resection procedures):
 - i. Greater than 10 degrees IMA OR Greater than 15 degrees MTP angle

DOCUMENTATION REQUIREMENTS: Molina Healthcare reserves the right to require that additional documentation be made available as part of its coverage determination; quality improvement; and fraud; waste and abuse prevention processes. Documentation required may include, but is not limited to, patient records, test results and credentials of the provider ordering or performing a drug or service. Molina Healthcare may deny reimbursement or take additional appropriate action if the documentation provided does not support the initial determination that the drugs or services were medically necessary, not investigational or experimental, and otherwise within the scope of benefits afforded to the member, and/or the documentation demonstrates a pattern of billing or other practice that is inappropriate or excessive.

SUMMARY OF MEDICAL EVIDENCE

Randomized Controlled Trials

Dragosloveanu et al. (2022) conducted a randomized controlled trial to evaluate open and minimally invasive surgical (MIS) technique approaches for chevron osteotomy. Forty patients were included in the study with 26 (24 female and 2 male) in the open chevron (OC) osteotomy (group and 24 (24 female and 0 male) in the MIS group. Patient were eligible for the study if they were older than 20 years, previously failed conservative treatment, and had a moderate

Molina Clinical Policy
Foot Surgery: Bunionectomy
Policy No. 700

Last Approval: 02/11/2026

Next Review Due By: February 2027



valgus deformity. Exclusion criteria included patients with previous first metatarsal osteotomy, instability of the first metatarsocuneiform, metatarsophalangeal joint osteoarthritis, and systemic disease affecting the musculoskeletal system. Outcomes included pain measured by the Visual Analog Scale (VAS), function measured with The American Orthopaedic Foot and Ankle Surgery score (AOFAS), hallux angle (HVA), and intramedullary angle (IMA). VAS scores were significantly better for the MIS group at discharge ($p < 0.001$), 3 weeks ($p < 0.001$), 6 weeks ($p < 0.001$), and 6 months ($p = 0.004$) post-surgery. At 6-months postoperative the VAS score was comparable between both groups ($p = 0.285$). There was no significant difference in AOFAS scores reported between groups pre- or post-operative ($p = 0.882$). While both groups demonstrated improvements in IMA and HVA at the last post-operative follow-up there was no significant differences reported between the groups. Complications reported included the screw prominence (3 MIS and 1 OS patient) and one metatarsalgia in the OC group. The study was limited by the small sample and short-term follow-up. The results demonstrate that minimally invasive and open techniques for chevron osteotomy have comparable results although more studies are needed to further evaluate the procedures.

Systematic Reviews and Meta-Analyses

Lewis et al. (2024) conducted a systematic review and meta-analysis to assess the clinical and radiographic outcomes of percutaneous or minimally invasive surgery for bunionette deformity correction. The review included 18 studies involving 714 feet in 580 patients, focusing on specific osteotomy techniques. Fourteen studies utilized unfixed distal osteotomy, while the remaining four employed K-wire fixation to maintain reduction. The analysis revealed statistically significant improvements in clinical outcomes, including the American Orthopaedic Foot & Ankle Society (AOFAS) scores and pain levels, as well as radiologic outcomes, such as the fourth-fifth intermetatarsal angle and fifth metatarsophalangeal angle. Complication rates ranged from 0% to 21.4%, with nonunion rates between 0% and 5.6%. The most common complication was hypertrophic callus formation, which typically resolved without requiring additional surgical intervention. Limitations of the studies included variability in case series, a lack of comparative research, and concurrent procedures performed with bunionette correction. Overall, the findings indicate that percutaneous techniques are safe and effective, offering significant improvements in radiographic alignment and patient-reported outcomes.

Peng et al. (2024) conducted a systematic review and meta-analysis to compare the outcomes of chevron and scarf osteotomies for correcting hallux valgus. The analysis included six randomized controlled trials involving 507 feet, with 261 treated using chevron osteotomy and 246 with scarf osteotomy. Primary outcomes measured were the hallux valgus angle (HVA), intermetatarsal angle (IMA), American Orthopedic Foot and Ankle Society (AOFAS) scores, and complication rates. The findings showed that chevron osteotomy resulted in significantly smaller postoperative HVAs compared to scarf osteotomy ($P < .00001$). However, differences in postoperative IMAs ($P = .19$), AOFAS scores ($P = .81$), and complication rates ($P = .53$) between the two procedures were not statistically significant. Limitations included the small number of RCTs analyzed, the limited use of AOFAS scores in only three studies, and small sample sizes. Overall, chevron osteotomy demonstrated better postoperative HVA correction, while the other outcomes showed no significant differences between the two techniques.

Nair et al. (2022) conducted a systematic review to evaluate open and minimally invasive surgical (MIS) techniques used for treating recurrent hallux valgus (HV). Ten articles including 273 patients, and 301 feet were included in the review. Articles included were published in the English language, included patient who had undergone revision surgery following primary HV correction surgery, and reported data points for primary and revision surgery. Review articles, expert opinion cases, and single case reports were excluded. Eighty-six (28.6%) patients underwent revision with MIS techniques with distal first metatarsal osteotomy (32) and modified subcapital metatarsal osteotomy (54). Open revision techniques included 80 scarf osteotomies (26.6%), 51 Lapidus procedures (16.9%), and 84 first MTP joint arthrodesis (27.9%). The American Orthopaedic Foot and Ankle Society (AOFAS) score, pain scoring, patient satisfaction, HV angle (HVA), and intermetatarsal angle (IMA) measurements were used. AOFAS score was an average of 38.3 for MIS revisions and 26.8 for traditional open techniques. MIS techniques had median postoperative reduction of 5.6 degrees IMA and 18.4 degrees HVA. Open revision techniques had a median of 4.4 degrees IMA and 18.4 degrees HVA. Pain scoring and patient satisfaction varied between studies which prevented the pooling of outcomes. Complications were reported in 74 surgeries and included nonunion, painful or broken metalwork requiring correction/removal, infection requiring antibiotics, transfer metatarsalgia, recurrent HV after revision, delayed union, malunion, and stiffness. While MIS techniques did not show worse outcomes or safety concerns compared to open techniques, due to inability to pool outcomes and directly compare studies this review was unable to make firm recommendation for MIS over open techniques.

Hernández-Castillejo et al. (2020) conducted a systematic review and analysis of 12 articles to assess the

Molina Clinical Policy
Foot Surgery: Bunionectomy
Policy No. 700

Last Approval: 02/11/2026

Next Review Due By: February 2027



effectiveness of hallux valgus surgery on patient quality of life (QoL). Inclusion criteria included patients > 16 years old, open or minimally invasive surgery procedures, QoL reports on physical, mental and social domains, included randomized controlled trials (RCTs), non-randomized experimental studies, and single-arm pre-post studies, and written in English or Spanish. Studies were excluded if they reported on foot or ankle pathologies other than hallux valgus (HV) and patients who had undergone HV revision surgery. Pooled effect size (ES) for QoL physical domain scores were 1.01 for body pain and 0.43 for physical function. Mental domain ES was 0.24 and 0.42 for social domain. The pooled difference in means for Vas score was -4.1. This study was limited by selection bias and limited availability of complete information from study reports. It was also noted that several clinician- and patient-related factors including surgical approach, surgeon skills, and comorbidities may have influenced outcomes. This review concluded that HV surgery results in decreased body pain, improved physical function, and improved social QoL, but did not affect mental QoL.

Barg et al. (2018) conducted a systematic review to analyze the outcomes of surgical correction for hallux valgus deformity. A total of 229 studies were included in the review with 16,273 procedures on 12,866 patients. Nine different surgery types were reviewed including distal osteotomy, proximal osteotomy, shaft osteotomy, joint hemiresection, simple bunionectomy, shaft and akin osteotomies, proximal and akin osteotomies, and first TMT arthrodesis. Outcomes measured included postoperative patient satisfaction, postoperative pain, recurrence of HV deformity, intraoperative nerve injury, rate of infection, osseous nonunion, rate of reoperation, and postoperative hallux varus deformity. A significant difference was reported in patient satisfaction across the nine different surgery types ($p = 0.004$). Dissatisfaction was highest in patient who underwent a simple bunionectomy or joint hemiresection. A significant difference was reported in the prevalence of postoperative metatarsalgia ($p < 0.001$) with the highest rates in patients who underwent joint hemiresection. Postoperative pain was comparable between the surgery types ($p = 0.026$). The rate of recurrent HV deformity ($p = 0.43$) and intraoperative nerve injury ($p = 0.38$) was comparable across surgery types. There was a significant difference in postoperative infection rates across the surgeries ($p = 0.02$). A significant difference in postoperative rates of osseous nonunion was reported across the different surgery types ($p = 0.012$). Patients who underwent a first tarsometatarsal arthrodesis had the highest rates of postoperative infection and osseous nonunion. Rate of reoperation was comparable across all surgery types ($p = 0.005$) as was the need for hardware removal ($p < 0.001$). Rates of postoperative hallux deformity was highest in patients who underwent a proximal osteotomy. It should be noted that the quality of the source literature, which is dominated by single-surgeon retrospective case series, as well as the extended period covered, with varied data-quality standards, limited the scope of this review. The systematic review found that arthrodesis and arthroplasty procedures had greater percentages of unfavorable outcomes.

National and Specialty Organizations

The **American College of Foot and Ankle Surgeons (ACFAS)** issued a clinical consensus statement “on a broad range of topics relevant to the clinical practice of foot and ankle surgeons as it relates to the hallux valgus deformity utilizing not only the best available evidence but also a degree of our clinical experience and common sense” (Meyr et al. 2022). Thirteen statements for evaluations considerations, perioperative considerations, and postoperative considerations were addressed and rated on an appropriateness scale* and includes the following:

- Radiographic evaluation is required for effective assessment of hallux valgus deformity (ACFAS Appropriate). Radiographic measurements help describe the severity of the HV. The most used parameters evaluate the alignment of the first and second metatarsals, the hallux, and the sesamoid complex in a transverse plane. A normal IMA is generally less than 8-10 degrees, and a normal HVA is typically less than 15-20 degrees. Increasing angular values correlate with greater deformity severity.
- The outcome of surgical correction is independent of procedure selection (ACFAS Appropriate)
 - No consensus reached on the concept of basing procedure selection on severity of deformity (ACFAS Neither appropriate nor inappropriate)
 - There may be multiple procedures that can achieve the goals of operative management in a single patient
- Use of biologic augmentation, such as bone marrow aspirate concentrate, may be considered in the surgical correction of hallux valgus (ACFAS Neither appropriate nor inappropriate)

* ACFAS appropriateness scale to attain consensus on clinical questions by members of the panel: Rating: 1, 2, 3, 4, 5, 6, 7, 8, 9. Inappropriate: 1, 2, or 3; Neither inappropriate or appropriate: 4, 5, or 6; Appropriate: 7, 8, or 9

CODING & BILLING INFORMATION

CPT (Current Procedural Terminology)

Code	Description
Bunionette	
28110	Ostectomy, partial excision, fifth metatarsal head (bunionette) (separate procedure)
28113	Ostectomy, complete excision; fifth metatarsal head
28307	Osteotomy, with or without lengthening, shortening or angular correction, metatarsal; first metatarsal with autograft (other than first toe)
28308	Osteotomy, with or without lengthening, shortening or angular correction, metatarsal; other than first metatarsal, each
Bunionectomy	
28292	Correction, hallux valgus with bunionectomy, with sesamoidectomy, when performed; with resection of proximal phalanx base, when performed, any method
28295	Correction, hallux valgus with bunionectomy, with sesamoidectomy, when performed; with proximal metatarsal osteotomy, any method
28296	Correction, hallux valgus with bunionectomy, with sesamoidectomy, when performed; with distal metatarsal osteotomy, any method
28297	Correction, hallux valgus with bunionectomy, with sesamoidectomy, when performed; with first metatarsal and medial cuneiform joint arthrodesis, any method
28298	Correction, hallux valgus with bunionectomy, with sesamoidectomy, when performed; with proximal phalanx osteotomy, any method
28299	Correction, hallux valgus with bunionectomy, with sesamoidectomy, when performed; with double osteotomy, any method
28306	Osteotomy, with or without lengthening, shortening or angular correction, metatarsal; first metatarsal
28310	Osteotomy, shortening, angular or rotational correction; proximal phalanx, first toe (separate procedure)
28312	Osteotomy, shortening, angular or rotational correction; other phalanges, any toe

CODING DISCLAIMER: Codes listed in this policy are for reference purposes only and may not be all-inclusive. Deleted codes and codes which are not effective at the time the service is rendered may not be eligible for reimbursement. Listing of a service or device code in this policy does not guarantee coverage. Coverage is determined by the benefit document. Molina adheres to Current Procedural Terminology (CPT®), a registered trademark of the American Medical Association (AMA). All CPT codes and descriptions are copyrighted by the AMA; this information is included for informational purposes only. Providers and facilities are expected to utilize industry standard coding practices for all submissions. When improper billing and coding is not followed, Molina has the right to reject/deny the claim and recover claim payment(s). Due to changing industry practices, Molina reserves the right to revise this policy as needed.

APPROVAL HISTORY

02/11/2026	Policy revised. Replaced list of required conservative treatments with broader language to allow for provider directed management. Removed general contraindications. Updated bunionette deformity angle criteria to require greater than 10 degrees IMA angle. Updated Summary of Medical Evidence and References. IRO Peer Review on February 10, 2026, by a practicing physician board certified in Orthopedic Surgery.
02/12/2025	Policy reviewed. Updated Summary of Medical Evidence and References. IRO Peer Review on December 13, 2024, by a practicing physician board-certified in Orthopedic Surgery.
12/11/2024	Clarified clinical indications by reorganizing criteria into one coverage section for all surgeries and removing E/I/U indications. Annual review will be completed as scheduled in 2025.
04/10/2024	Policy reviewed, no changes to criteria. Updated Summary of Medical Evidence and References.
04/13/2023	New policy, replaces MCP-401: Foot Surgery. IRO Peer Review March 31, 2023, by a practicing, board-certified physician in Orthopedic Surgery.

REFERENCES

- Barg A., Harmer JR, Presson AP, et al. Unfavorable outcomes following surgical treatment of hallux valgus deformity: A systematic literature review. J Bone Joint Surg Am. 2018 Sep 19;100(18):1563-1573. doi: 10.2106/JBJS.17.00975.
- Dragosloveanu S, Popov VM, Cotor DC, Dragosloveanu C, Stoica CI. Percutaneous Chevron Osteotomy: A prospective randomized controlled trial. Medicina (Kaunas). 2022 Mar 1;58(3):359. doi: 10.3390/medicina58030359. PMID: 35334535; PMCID: PMC8948867

Molina Clinical Policy
Foot Surgery: Bunionectomy
Policy No. 700

Last Approval: 02/11/2026

Next Review Due By: February 2027



3. DynaMed. EBSCO Information Services. Management of hallux valgus (bunion). Updated February 6, 2025. Accessed January 12, 2026. <http://www.dynamed.com>.
4. Ettinger S, Spindler FT, Marschall U, et al. Hallux Valgus: Prevalence and Treatment Options. *Dtsch Arztebl Int*. 2025 May 30;122(11):308-314. doi: 10.3238/arztebl.m2025.0068. PMID: 40332014; PMCID: PMC12539686.
5. Ferrari J. & Monson E. Hallux valgus deformity (bunion) in adults. Updated May 20, 2025. Accessed January 13, 2026. <http://www.uptodate.com>.
6. Hernández-Castillejo LE, Martínez Vizcaíno V, Garrido-Miguel M, et al. Effectiveness of hallux valgus surgery on patient quality of life: a systematic review and meta-analysis. *Acta Orthop*. 2020 Aug;91(4):450-456. doi: 10.1080/17453674.2020.1764193. Epub 2020 May 14. PMID: 32408787; PMCID: PMC8023907.
7. Kuhn J, Alvi F. Hallux Valgus. In: StatPearls [Internet]. Treasure Island (FL). Updated August 28, 2023. Accessed January 13, 2026. <https://www.ncbi.nlm.nih.gov/books/NBK553092/>.
8. Lewis TL, Lam P, Alkhalafan Y, et al. Minimally invasive surgery for management of bunionette deformity (Tailor's Bunion) using fifth metatarsal osteotomies: A systematic review and meta-analysis. *Foot Ankle Orthop*. 2024 Jul 29;9(3):24730114241263095. doi: 10.1177/24730114241263095. PMID: 39086382; PMCID: PMC11289809.
9. Meyr AJ, Doyle MD, King CM, et al. The American College of Foot and Ankle Surgeons clinical consensus statement: Hallux valgus. *J Foot Ankle Surg*. 2022 Mar-Apr;61(2):369-383. doi: 10.1053/j.jfas.2021.08.011. Epub 2021 Sep 4. PMID: 34706857.
10. Nair A, Bence M, Saleem J, et al. A systematic review of open and minimally invasive surgery for treating recurrent hallux valgus. *Surg J (N Y)*. 2022 Dec 21;8(4):e350-e356. doi: 10.1055/s-0042-1759812. PMID: 36568477; PMCID: PMC9771687.
11. Peng YN, Peng YH, & Chen CPC. Chevron osteotomy and scarf osteotomy for hallux valgus angle and intermetatarsal angle correction: A systematic review and meta-analysis of randomized controlled trials. *J Orthop Surg Res*. 2024 Sep 14;19(1):566. doi: 10.1186/s13018-024-05007-0. PMID: 39272201; PMCID: PMC11401393.
12. Ray JJ, Friedmann AJ, Hanselman AE, et al. Hallux valgus. *Foot Ankle Orthop*. 2019 May 7;4(2):2473011419838500. doi: 10.1177/2473011419838500.
13. Shi GG, Humayun A, Whalen JL, et al. Management of bunionette deformity. *J Am Acad Orthop Surg*. 2018 Oct 1. 26 (19):e396-e404. doi: 10.5435/JAAOS-D-17-00345.

APPENDIX

Reserved for State specific information. Information includes, but is not limited to, State contract language, Medicaid criteria and other mandated criteria.

Washington

For Medicaid, there is a language around foot care in the following Washington Administrative Codes: WAC 182-531-0150: Noncovered physician-related and health care professional services—General and administrative; and WAC 182-531-1300: Foot care services for clients twenty-one years of age and older. Per the WACs, routine foot care is considered a non-covered healthcare service unless the client meets conditions and criteria outlined in WAC 182-531-1300. If criteria are needed for medical necessity review in the case that the member has a qualifying condition, this MCP can be applied as a medical necessity tool.