

# Why Are Immunizations So Important?

Immunizations are shots that help the body fight diseases. Each shot fights a different disease like chickenpox, measles, or mumps. Kids need shots to fight disease and to be ready for school.

Talk to a doctor about the shots your kid needs to be protected from diseases. **Remember, these shots are covered at no cost to you.**

Here's a look at the shot schedule recommended by the Centers for Disease Control and Prevention (CDC):



## BIRTH

Hepatitis B (HepB)



## 1-2 MONTHS

HepB (2<sup>nd</sup> Dose)



## 2 MONTHS

Rotavirus (RV)  
Diphtheria, Tetanus,  
Acellular Pertussis (DTaP)  
Haemophilus Influenza  
Type B (Hib)  
Pneumococcal  
Conjugate (PCV13)  
Inactivated  
Poliovirus (IPV)



## 4 MONTHS

RV (2<sup>nd</sup> Dose)  
DTaP (2<sup>nd</sup> Dose)  
Hib (2<sup>nd</sup> Dose)  
PCV13 (2<sup>nd</sup> Dose)  
IPV (2<sup>nd</sup> Dose)



## 6 MONTHS

HepB (3<sup>rd</sup> Dose 6-18 months)  
RV (3<sup>rd</sup> Dose)  
DTaP (3<sup>rd</sup> Dose)  
Hib (3<sup>rd</sup> Dose)  
PCV13 (3<sup>rd</sup> Dose)  
IPV (3<sup>rd</sup> Dose 6-18 months)



## 12-15 MONTHS

Hib (4<sup>th</sup> Dose)  
PCV13 (4<sup>th</sup> Dose)  
Measles, Mumps,  
Rubella (MMR)  
Varicella (VAR)  
Hepatitis A (HepA)



## 15-18 MONTHS

DTaP (4<sup>th</sup> Dose)



## 18 MONTHS

HepA (2<sup>nd</sup> Dose)  
Total of 2 doses:  
12-23 months



**Influenza (Flu Shot) Yearly: 6 months to 18 years**  
**Some kids under 8 may get 2 flu shots every year**

Doctors may choose different types of vaccines that may have different dosages. So your child may or may not get an extra dose of some immunizations. Talk to your doctor for more information.

Source: Centers for Disease Control and Prevention (CDC)

300900THMDKYEN 221210



**2 YEARS**



**4-6 YEARS**

DTaP (5<sup>th</sup> Dose)  
IPV (4<sup>th</sup> Dose)  
MMR (2<sup>nd</sup> Dose)  
VAR (2<sup>nd</sup> Dose)



**11-12 YEARS**

Tetanus, Diphtheria,  
Acellular Pertussis  
(Tdap)  
Meningococcal  
Human  
Papillomavirus  
(HPV) 2 doses



Most children will get their HPV shot at age 11. Some children with certain risk factors can get the HPV shot as early as age 9. Some children may need a 3rd dose. Talk to your doctor to find out what is best for your child.



**16 YEARS**

Meningococcal



**Influenza (Flu Shot) Yearly: 6 months to 18 years**  
**Some kids under 8 may get 2 flu shots every year**



MARK04782 APP\_X/XX/2021