Molina Healthcare of Texas

Medicaid HEDIS® Annual Rates: 2018–2020

Molina's scores for the past three years are below. You can see how Molina performs compared to our target goal.

| Measure/Data Element | 2018 | 2019 | 2020 | 2020 Goal |
|---|---|---|---|---|
| Adherence to Antipsychotic Medications for Individuals with Schizophrenia (SAA) | 52.97% | 54.30% | 55.47% | 67.47% |
| Adult BMI Assessment (ABA) | 87.31% | 81.51% | 90.38% | 93.67% |
| Annual Dental Visits (ADV) | NB | NB | NB | 64.11% |
| Antidepressant Medication Management (AMM) Effective Continuation Phase Treatment | 35.33% | 35.64% | 36.66% | 40.95% |
| Appropriate Testing for Pharyngitis (CWP) | 73.11% | 77.70% | 79.24% | 86.51% |
| Appropriate Treatment for Upper Respiratory Infection (URI) | 85.58% | 90.40% | 83.81% | 94.88% |
| Asthma Medication Ratio (AMR) Total | 55.06% | 58.08% | 55.73% | 68.52% |
| Avoidance of Antibiotic Treatment for Acute Bronchitis / Bronchiolitis (AAB) | 26.35% | 31.04% | 57.89% | 41.07% |
| Breast Cancer Screening (BCS) | 54.60% | 55.85% | 56.75% | 63.98% |
| Cervical Cancer Screening (CCS) | 55.72% | 52.07% | 52.07% | 66.49% |
| Childhood Immunization Status (CIS) Combination #10 | 27.98% | 29.20% | 29.20% | 42.02% |
| Chlamydia Screening in Women (CHL) Total | 44.09% | 47.55% | 43.91% | 66.24% |
| Comprehensive Diabetes Care (CDC) BP Control < 140/90 | 45.50% | 54.01% | 54.01% | 71.44% |
| Comprehensive Diabetes Care (CDC) HbA1c Control < 8.0% | 39.42% | 44.77% | 44.77% | 55.96% |
| Comprehensive Diabetes Care (CDC) Eye Exam | 59.85% | 61.80% | 61.80% | 64.72% |
| Controlling High Blood Pressure (CBP) | 43.80% | 52.80% | 52.80% | 66.91% |
| Diabetes Screening for People with Schizophrenia or Bipolar Disorder Who Are Using Antipsychotic Medications (SSD) | 81.96% | 84.13% | 84.09% | 84.27% |
| Follow-Up After Emergency Department Visit for Alcohol and Other Drug Abuse or Dependence (FUA) 7-Day Follow-Up Total | 4.27% | 4.33% | 2.57% | 16.93% |
| Follow-Up After Emergency Department Visit for Mental Illness (FUM) 7-Day Follow-Up Total | 32.10% | 29.21% | 28.78% | 49.65% |
| Follow-Up After Hospitalization for Mental Illness (FUH) 7-Day Follow-Up Total | 33.96% | 37.86% | 37.23% | 42.69% |
| Follow-Up Care for Children Prescribed ADHD Medication (ADD) Continuation and Maintenance Phase | 69.59% | 70.00% | 56.76% | 62.69% |
| Immunizations for Adolescents (IMA) Combination #2 | 42.82% | 42.58% | 46.96% | 40.39% |

NA = Not Available Due to Small Denominator        NQ = Not Required

NB = No Benefit        BR = Biased Rate

NR = Not Reported

* Reported rate calculated per 1000 members

| Measure/Data Element | 2018 | 2019 | 2020 | 2020 Goal |
|---|---|---|---|---|
| Initiation and Engagement of Alcohol and Other Drug Dependence Treatment (IET) Engagement of AOD Treatment Total | 6.05% | 6.31% | 5.96% | 18.12% |
| Medication Management for People with Asthma (MMA) Medication Compliance 75% Total | 30.02% | 31.90% | 29.60% | 42.77% |
| Metabolic Monitoring for Children and Adolescents on Antipsychotics (APM) Total | 28.61% | 33.16% | 34.12% | 40.90% |
| Pharmacotherapy Management of COPD Exacerbation (PCE) Systemic Corticosteroid | 69.52% | 67.83% | 70.16% | 75.38% |
| Pharmacotherapy Management of COPD Exacerbation (PCE) Bronchodilator | 85.81% | 86.71% | 84.64% | 87.93% |
| Prenatal and Postpartum Care (PPC) Timeliness of Prenatal Care | 85.06% | 86.62% | 98.54% | 87.59% |
| Prenatal and Postpartum Care (PPC) Postpartum Care | 54.68% | 56.69% | 79.32% | 69.83% |
| Statin Therapy for Patients with Cardiovascular Disease (SPC) Received Statin Therapy Total | 73.12% | 75.83% | 79.71% | 81.24% |
| Statin Therapy for Patients with Cardiovascular Disease (SPC) Statin Adherence 80% Total | 55.14% | 57.06% | 59.36% | 69.97% |
| Statin Therapy for Patients with Diabetes (SPD) Received Statin Therapy | 67.22% | 69.31% | 69.97% | 67.19% |
| Statin Therapy for Patients with Diabetes (SPD) Statin Adherence 80% | 53.37% | 56.86% | 57.07% | 66.42% |
| Use of First-Line Psychosocial Care for Children and Adolescents on Antipsychotics (APP) Total | 48.51% | 45.07% | 48.69% | 66.58% |
| Use of Imaging Studies for Low Back Pain (LBP) | 65.56% | 65.87% | 73.17% | 76.02% |
| Use of Opioids at High Dosage (HDO) | 18.50* | 0.90% | 1.24% | ≤ 2.27% |
| Use of Opioids from Multiple Providers (UOP) Multiple Prescribers and Multiple Pharmacies | 22.02* | 2.07% | 1.32% | ≤ 2.14% |
| Weight Assessment and Counseling for Nutrition and Physical Activity for Children/Adolescents (WCC) BMI Percentile | 68.58% | 70.56% | 73.95% | 85.16% |

NA = Not Available Due to Small Denominator     NQ = Not Required

NB = No Benefit     BR = Biased Rate

NR = Not Reported

* Reported rate calculated per 1000 members