Molina Healthcare of Wisconsin

Medicaid HEDIS® Annual Rates: 2018–2020

Molina's scores for the past three years are below. You can see how Molina performs compared to our target goal.

| Measure/Data Element | 2018 | 2019 | 2020 | 2020 Goal |
|---|---|---|---|---|
| Adherence to Antipsychotic Medications for Individuals with Schizophrenia (SAA) | 45.83% | 39.47% | 38.71% | 67.47% |
| Adult BMI Assessment (ABA) | 92.75% | 91.73% | 92.52% | 93.67% |
| Annual Dental Visits (ADV) | 57.85% | 57.42% | 58.06% | 64.11% |
| Antidepressant Medication Management (AMM) Effective Continuation Phase Treatment | 35.40% | 31.62% | 31.18% | 40.95% |
| Appropriate Testing for Pharyngitis (CWP) | 81.18% | 86.34% | 82.99% | 86.51% |
| Appropriate Treatment for Upper Respiratory Infection (URI) | 85.32% | 90.27% | 89.45% | 94.88% |
| Asthma Medication Ratio (AMR) | 45.28% | 54.33% | 56.38% | 68.52% |
| Avoidance of Antibiotic Treatment for Acute Bronchitis / Bronchiolitis (AAB) | 34.29% | 36.76% | 58.14% | 41.07% |
| Breast Cancer Screening (BCS) | 64.63% | 63.43% | 59.92% | 63.98% |
| Cervical Cancer Screening (CCS) | 70.45% | 70.80% | 72.02% | 66.49% |
| Childhood Immunization Status (CIS) Combination #10 | 36.24% | 34.31% | 34.06% | 42.02% |
| Chlamydia Screening in Women (CHL) Total | 63.97% | 64.67% | 63.22% | 66.24% |
| Comprehensive Diabetes Care (CDC) BP Control < 140/90 | 64.23% | 69.83% | 69.83% | 71.44% |
| Comprehensive Diabetes Care (CDC) HbA1c Control < 8.0% | 46.47% | 54.50% | 54.50% | 55.96% |
| Comprehensive Diabetes Care (CDC) Eye Exam | 46.96% | 49.39% | 49.39% | 64.72% |
| Controlling High Blood Pressure (CBP) | 54.88% | 51.82% | 55.23% | 66.91% |
| Diabetes Screening for People with Schizophrenia or Bipolar Disorder Who Are Using Antipsychotic Medications (SSD) | 81.71% | 78.93% | 74.67% | 84.27% |
| Follow-Up After Emergency Department Visit for Alcohol and Other Drug Abuse or Dependence (FUA) 7-Day Follow-Up Total | 9.23% | 11.73% | 11.45% | 16.93% |
| Follow-Up After Emergency Department Visit for Mental Illness (FUM) 7-Day Follow-Up Total | 32.99% | 35.60% | 42.61% | 49.65% |
| Follow-Up After Hospitalization for Mental Illness (FUH) 7-Day Follow-Up Total | 41.56% | 35.35% | 44.85% | 42.69% |
| Follow-Up for Children Prescribed ADHD Medication (ADD) Continuation and Maintenance Phase | 63.64% | 50.00% | 47.73% | 62.69% |

NA = Not Available Due to Small Denominator          NQ = Not Required

NB = No Benefit          BR = Biased Rate

NR = Not Reported

\* Reported rate calculated per 1000 members

| Measure/Data Element | 2018 | 2019 | 2020 | 2020 Goal |
|---|---|---|---|---|
| Immunizations for Adolescents (IMA) Combination #2 | 37.71% | 42.58% | 42.58% | 40.39% |
| Initiation and Engagement of Alcohol and Other Drug Dependence Treatment (IET) Engagement of AOD Treatment Total | 12.46% | 14.40% | 17.36% | 18.12% |
| Medication Management for People with Asthma (MMA) Medication Compliance 75% Total | 23.11% | 24.31% | 30.53% | 42.77% |
| Metabolic Monitoring for Children and Adolescents on Antipsychotics (APM) Total | 11.69% | 8.33% | 19.05% | 40.90% |
| Pharmacotherapy Management of COPD Exacerbation (PCE) Systemic Corticosteroids | 81.94% | 72.41% | 82.86% | 75.38% |
| Pharmacotherapy Management of COPD Exacerbation (PCE) Bronchodilator | 87.50% | 87.93% | 88.57% | 87.93% |
| Prenatal and Postpartum Care (PPC) Timeliness of Prenatal Care | 81.44% | 79.81% | 92.70% | 87.59% |
| Prenatal and Postpartum Care (PPC) Postpartum Care | 56.70% | 59.37% | 71.29% | 69.83% |
| Statin Therapy for Patients with Cardiovascular Disease (SPC) Received Statin Therapy Total | 85.33% | 84.51% | 83.82% | 81.24% |
| Statin Therapy for Patients with Cardiovascular Disease (SPC) Statin Adherence 80% Total | 68.75% | 63.33% | 64.91% | 69.97% |
| Statin Therapy for Patients with Diabetes (SPD) Received Statin Therapy | 65.96% | 65.38% | 67.05% | 67.19% |
| Statin Therapy for Patients with Diabetes (SPD) Statin Adherence 80% | 54.44% | 62.61% | 62.29% | 66.42% |
| Use of First-Line Psychosocial Care for Children and Adolescents on Antipsychotics (APP) Total | 63.64% | 57.50% | 57.14% | 66.58% |
| Use of Imaging Studies for Low Back Pain (LBP) | 73.70% | 69.88% | 75.41% | 76.02% |
| Use of Opioids at High Dosage (HDO) | 185.42* | NQ | 25.97% | ≤ 2.27% |
| Use of Opioids from Multiple Providers (UOP) Multiple Prescribers and Multiple Pharmacies | 121.47* | NQ | 2.98% | ≤ 2.14% |
| Weight Assessment and Counseling for Nutrition and Physical Activity for Children/Adolescents (WCC) BMI Percentile | 73.24% | 80.29% | 75.61% | 85.16% |

NA = Not Available Due to Small Denominator          NQ = Not Required
NB = No Benefit                                       BR = Biased Rate
NR = Not Reported

* Reported rate calculated per 1000 members