



Molina Healthcare of Florida

Medicaid HEDIS® Annual Rates: 2018-2020

Molina’s scores for the past three years are below. You can see how Molina performs compared to our target goal.

| Measure/Data Element | 2018 | 2019 | 2020 | 2020 Goal |
|---|--------|--------|--------|-----------|
| Adherence to Antipsychotic Medications for Individuals with Schizophrenia (SAA) | 57.54% | 60.95% | 64.42% | 67.47% |
| Adult BMI Assessment (ABA) | 88.21% | 90.75% | 96.86% | 93.67% |
| Annual Dental Visits (ADV) Total | 50.06% | 48.48% | NB | 64.11% |
| Antidepressant Medication Management (AMM) Effective Continuation Phase Treatment | 36.02% | 39.37% | 41.53% | 40.95% |
| Appropriate Testing for Pharyngitis (CWP) | 72.98% | 70.09% | 70.10% | 86.51% |
| Appropriate Treatment for Upper Respiratory Infection (URI) | 89.51% | 90.18% | 87.53% | 94.88% |
| Asthma Medication Ratio (AMR) Total | 63.95% | 71.33% | 81.82% | 68.52% |
| Avoidance of Antibiotic Treatment for Acute Bronchitis / Bronchiolitis (AAB) | 28.16% | 31.84% | 57.97% | 41.07% |
| Breast Cancer Screening (BCS) | 65.18% | 60.91% | 69.21% | 63.98% |
| Cervical Cancer Screening (CCS) | 63.99% | 56.93% | 63.26% | 66.49% |
| Childhood Immunization Status (CIS) Combination #10 | 27.25% | 24.82% | 38.05% | 42.02% |
| Chlamydia Screening in Women (CHL) Total | 63.90% | 64.67% | 67.81% | 66.24% |
| Comprehensive Diabetes Care (CDC) BP Control < 140/90 | 52.07% | 54.26% | 71.05% | 71.44% |
| Comprehensive Diabetes Care (CDC) HbA1c Control < 8.0% | 48.91% | 40.88% | 56.93% | 55.96% |
| Comprehensive Diabetes Care (CDC) Eye Exam | 58.15% | 58.15% | 65.21% | 64.72% |
| Controlling High Blood Pressure (CBP) | 50.36% | 54.74% | 72.13% | 66.91% |
| Diabetes Screening for People with Schizophrenia or Bipolar Disorder Who Are Using Antipsychotic Medications (SSD) | 82.89% | 83.06% | 85.74% | 84.27% |
| Follow-Up After Emergency Department Visit for Alcohol and Other Drug Abuse or Dependence (FUA) 7-Day Follow-Up Total | 3.73% | 5.60% | 5.92% | 16.93% |
| Follow-Up After Emergency Department Visit for Mental Illness (FUM) 7-Day Follow-Up Total | 21.44% | 25.81% | 22.39% | 49.65% |
| Follow-Up After Hospitalization for Mental Illness (FUH) 7-Day Follow-Up Total | 28.54% | 31.53% | 38.83% | 42.69% |
| Follow-Up for Children Prescribed ADHD Medication (ADD) Continuation and Maintenance Phase | 60.47% | 71.07% | 62.59% | 62.69% |

NA = Not Available Due to Small Denominator

NB = No Benefit

NR = Not Reported

*Reported rate calculated per 1000 members

NQ = Not Required

BR = Biased Rate

| Measure/Data Element | 2018 | 2019 | 2020 | 2020 Goal |
|---|--------|--------|--------|-----------|
| Immunizations for Adolescents (IMA) Combination #2 | 28.71% | 32.85% | 43.41% | 40.39% |
| Initiation and Engagement of Alcohol and Other Drug Dependence Treatment (IET) Engagement of AOD Treatment Total | 6.55% | 6.70% | 7.45% | 18.12% |
| Medication Management for People with Asthma (MMA) Medication Compliance 75% Total | 29.05% | 33.90% | 32.79% | 42.77% |
| Metabolic Monitoring for Children and Adolescents on Antipsychotics (APM) Total | 39.06% | 40.58% | 44.55% | 40.90% |
| Pharmacotherapy Management of COPD Exacerbation (PCE) Systemic Corticosteroids | 69.05% | 69.95% | 68.53% | 75.38% |
| Pharmacotherapy Management of COPD Exacerbation (PCE) Bronchodilator | 88.64% | 85.57% | 91.37% | 87.93% |
| Prenatal and Postpartum Care (PPC) Timeliness of Prenatal Care | 84.05% | 85.16% | 99.03% | 87.59% |
| Prenatal and Postpartum Care (PPC) Postpartum Care | 67.09% | 65.94% | 77.13% | 69.83% |
| Statin Therapy for Patients with Cardiovascular Disease (SPC) Received Statin Therapy Total | 71.99% | 77.79% | 79.77% | 81.24% |
| Statin Therapy for Patients with Cardiovascular Disease (SPC) Statin Adherence 80% Total | 58.29% | 66.73% | 64.59% | 69.97% |
| Statin Therapy for Patients with Diabetes (SPD) Received Statin Therapy | 62.70% | 67.42% | 76.06% | 67.19% |
| Statin Therapy for Patients with Diabetes (SPD) Statin Adherence 80% | 53.67% | 63.28% | 60.66% | 66.42% |
| Use of First-Line Psychosocial Care for Children and Adolescents on Antipsychotics (APP) Total | 62.63% | 64.99% | 77.39% | 66.58% |
| Use of Imaging Studies for Low Back Pain (LBP) | 61.53% | 63.25% | 66.41% | 76.02% |
| Use of Opioids at High Dosage (HDO) | 59.30* | 5.07% | 13.55% | ≤ 2.27% |
| Use of Opioids from Multiple Providers (UOP) Multiple Prescribers and Multiple Pharmacies | 51.01* | 3.76% | 2.04% | ≤ 2.14% |
| Weight Assessment and Counseling for Nutrition and Physical Activity for Children/Adolescents (WCC) BMI Percentile | 85.54% | 84.67% | 89.74% | 85.16% |

The reporting year is subsequent of the year reflected in the data; for example, “HEDIS 2020” is data collected from calendar year January-December 2019.

NA = Not Available Due to Small Denominator

NB = No Benefit

NR = Not Reported

*Reported rate calculated per 1000 members

NQ = Not Required

BR = Biased Rate