



Your Extended Family.

## Molina Healthcare of Michigan

### Medicaid HEDIS Annual Rates: 2016 – 2018

Molina's scores for the past three years are below. You can see how Molina performs compared to our target goal.

| Measure/Data Element  | 2016   | 2017   | 2018   | 2018 Goal |
|---|--------|--------|--------|-----------|
| Adult BMI Assessment (ABA)  | 90.15% | 97.14% | 96.00% | 92.46%    |
| Annual Dental Visits (ADV)  | NB     | 41.18% | 36.14% | 64.16%    |
| Antidepressant Medication Management (AMM)<br>Effective Acute Phase Treatment   | 51.46% | 48.20% | 54.54% | 57.82%    |
| Antidepressant Medication Management (AMM)<br>Effective Continuation Phase Treatment                                  | 34.29% | 32.61% | 37.54% | 42.31%    |
| Appropriate Testing for Children with Pharyngitis (CWP)   | 62.82% | 67.17% | 75.12% | 85.99%    |
| Appropriate Treatment for Children with Upper Respiratory<br>Infection (URI)  | 88.44% | 86.82% | 87.40% | 93.77%    |
| Medication Management for People With Asthma (MMA)<br>Total Medication Compliance 75% Rate only                       | 30.92% | 34.13% | 38.56% | 43.06%    |
| Asthma Medication Ratio (AMR)   | 61.35% | 60.91% | 63.06% | 67.03%    |
| Avoidance of Antibiotic Treatment in Adults With Acute<br>Bronchitis (AAB)  | 27.70% | 30.18% | 33.02% | 37.36%    |
| Breast Cancer Screening (BCS)   | 59.67% | 60.31% | 61.50% | 64.12%    |
| Cervical Cancer Screening (CCS)   | 65.63% | 65.69% | 72.34% | 66.01%    |
| Childhood Immunization Status (CIS)<br>Combination #10  | 32.23% | 30.24% | 31.31% | 40.88%    |
| Chlamydia Screening in Women (CHL)<br>Total Rate  | 66.63% | 66.23% | 67.35% | 65.43%    |
| Comprehensive Diabetes Care (CDC)<br>HbA1c Testing  | 86.04% | 87.64% | 90.42% | 90.45%    |
| Comprehensive Diabetes Care (CDC)<br>BP Control <140/90   | 55.41% | 55.19% | 51.11% | 70.76%    |
| Comprehensive Diabetes Care (CDC)<br>HbA1c Control <8.0%  | 50.90% | 56.73% | 54.55% | 55.47%    |
| Comprehensive Diabetes Care (CDC)<br>Eye Exam   | 57.43% | 62.03% | 62.16% | 64.23%    |
| Comprehensive Diabetes Care (CDC)<br>Monitoring for Nephropathy   | 92.12% | 90.73% | 92.87% | 92.05%    |
| Controlling High Blood Pressure (CBP)   | 53.60% | 49.04% | 51.82% | 65.78%    |
| Diabetes Screening for People with Schizophrenia or Bipolar<br>Disorder Who Are Using Antipsychotic Medications (SSD) | 84.61% | 83.10% | 85.87% | 84.78%    |
| Follow-Up After Hospitalization for Mental Illness (FUH)<br>7-Day Rate Only   | NQ     | NQ     | NB     | 45.79%    |

NA = Not Available Due to Small Denominator

NB = No Benefit

NR = Not Reported

NQ = Not Required

BR = Biased Rate

**Your Extended Family.**

| <b>Measure/Data Element</b>   | <b>2016</b> | <b>2017</b> | <b>2018</b> | <b>2018 Goal</b> |
|---|-------------|-------------|-------------|------------------|
| Follow-Up for Children Prescribed ADHD Medication (ADD)<br>Initiation Phase   | 37.42%      | 48.40%      | 48.91%      | 50.82%           |
| Follow-Up for Children Prescribed ADHD Medication (ADD)<br>Continuation Phase   | 45.83%      | 65.97%      | 37.54%      | 63.72%           |
| Human Papillomavirus Vaccine for Female Adolescents<br>(HPV)  | 24.94%      | 28.70%      | 33.17%      | 40.63%           |
| Immunizations for Adolescents (IMA)<br>Combination #1   | 90.54%      | 90.07%      | 86.87%      | 85.64%           |
| Initiation and Engagement of Alcohol and Other Drug<br>Dependence Treatment (IET)<br>Engagement of AOD Treatment Rate Total                     | 2.40%       | NQ          | NB          | 17.73%           |
| Medical Assistance With Smoking and Tobacco Use<br>Cessation (MSC)<br>Advising Smokers and Tobacco Users to Quit Rate only                      | --          | --          | --          | 80.87%           |
| Pharmacotherapy Management of COPD Exacerbation (PCE)<br>Systemic Corticosteroids   | 75.48%      | 73.20%      | 74.55%      | 76.27%           |
| Pharmacotherapy Management of COPD Exacerbation (PCE)<br>Bronchodilator   | 84.84%      | 87.74%      | 89.07%      | 87.60%           |
| Prenatal and Postpartum Care (PPC)<br>Timeliness of Prenatal Care   | 78.20%      | 83.33%      | 77.32%      | 87.06%           |
| Prenatal and Postpartum Care (PPC)<br>Postpartum Care   | 67.87%      | 75.80%      | 73.80%      | 69.34%           |
| Use of Imaging Studies for Low Back Pain (LBP)  | 77.61%      | 73.67%      | 72.99%      | 75.96%           |
| Weight Assessment and Counseling for Nutrition and Physical<br>Activity for Children/Adolescents (WCC)<br>BMI Screening Total                   | 80.46%      | 80.61%      | 84.64%      | 82.63%           |
| Weight Assessment and Counseling for Nutrition and Physical<br>Activity for Children/Adolescents (WCC)<br>Counseling for Nutrition              | 67.82%      | 71.39%      | 76.82%      | 77.91%           |
| Weight Assessment and Counseling for Nutrition and Physical<br>Activity for Children/Adolescents (WCC)<br>Counseling on Physical Activity Total | 63.68%      | 63.59%      | 68.75%      | 71.29%           |

NA = Not Available Due to Small Denominator

NB = No Benefit

NR = Not Reported

NQ = Not Required

BR = Biased Rate