

Request for Prior Authorization

FAX Completed Form To I (800) 574-2515

> Provider Help Desk I (877) 776-I567

ANTIDEPRESSANTS

(PLEASE PRINT – ACCURACY IS IMPORTANT)

IA Medicaid Member ID #	Patient name			DOB	
Patient address					
Provider NPI	Prescriber name			Phone	
Prescriber address	<u> </u>			Fax	
Pharmacy name	Address			Phone	
Prescriber must complete all informa Pharmacy NPI	ation above. It must be legible. Pharmacy fax		nplete or for	m will be returned.	
	,				
dosing, contraindications, warning Documentation of a previous trial 3) Documentation of a previous trial and 4) Documentation of a previous antidepressant; and 5) Document and 6) Documentation of a previous trial	I and therapy failure at a the rial and therapy failure at a ous trial and therapy failure cation of a previous trial and ous trial and therapy failure	erapeutic dose of therapeutic dos at a therapeutic therapy failure at a therapeutic	with two pr se with one c dose with at a therap c dose with	eferred generic SSI preferred generic Sone non-SSRI/SNR peutic dose with vila vortioxetine; and 7	RÍs; and SNRI; I generio azodone)
and 8) If the request is for dextromust include a previous trial and agent; and 9) If the request is for a must be with the preferred parendocumented intolerance. The requise of these agents would be med Non-Preferred Aplenzin Auvelity	methorphan and bupropion inadequate response at a th an isomer, prodrug or meta at drug of the same chemica quired trials may be override	extended- rele erapeutic dose bolite of the re I entity that res den when docur	ase tablet (with an ext quested me sulted in a p	Auvelity), one of the cended-release bup a cedication, one of the certial response with	e trials ropion e trials h a
must include a previous trial and agent; and 9) If the request is for a must be with the preferred paren documented intolerance. The requise of these agents would be med Non-Preferred Aplenzin Auvelity	methorphan and bupropion inadequate response at a th an isomer, prodrug or metal drug of the same chemical quired trials may be overrided lically contraindicated. Fetzima Other:	extended- rele erapeutic dose bolite of the re I entity that res den when docur	ase tablet (with an ext quested me sulted in a p mented evic	Auvelity), one of the ended-release bup edication, one of the eartial response with dence is provided the	e trials ropion e trials h a
must include a previous trial and agent; and 9) If the request is for must be with the preferred paren documented intolerance. The requise of these agents would be med Non-Preferred Aplenzin Auvelity Strength Dosage Instr	methorphan and bupropion inadequate response at a th an isomer, prodrug or metant drug of the same chemical puired trials may be overridually contraindicated.	extended- rele erapeutic dose bolite of the re I entity that res den when docur	ase tablet (with an ext quested me sulted in a p mented evic	Auvelity), one of the ended-release bup edication, one of the eartial response with dence is provided the	e trials ropion e trials h a
must include a previous trial and agent; and 9) If the request is for a must be with the preferred paren documented intolerance. The requise of these agents would be med Non-Preferred Aplenzin Auvelity	methorphan and bupropion inadequate response at a than isomer, prodrug or metal than isomer, prodrug trials may be override lically contraindicated. Fetzima Other:	extended- rele erapeutic dose bolite of the re I entity that res den when docur Quantity	ase tablet (with an ext quested me sulted in a p mented evid	Auvelity), one of the ended-release bupiedication, one of the edication, one of the edication artial response with dence is provided the epply	e trials ropion e trials h a
must include a previous trial and agent; and 9) If the request is for a must be with the preferred paren documented intolerance. The requise of these agents would be med Non-Preferred Aplenzin Auvelity Strength Dosage Instruction Diagnosis: Preferred Generic SSRI Trial I: D	methorphan and bupropion inadequate response at a the an isomer, prodrug or metal at drug of the same chemical quired trials may be override lically contraindicated. Fetzima Other:	extended- rele erapeutic dose bolite of the re I entity that res den when docur _ Quantity	ase tablet (with an ext quested me sulted in a p mented evic	Auvelity), one of the ended-release buping dication, one of the edication, one of the edication artial response with dence is provided the ended the edication are poly	e trials ropion e trials h a
must include a previous trial and agent; and 9) If the request is for must be with the preferred parent documented intolerance. The requise of these agents would be med Non-Preferred Aplenzin Auvelity Strength Dosage Instruction Diagnosis: Preferred Generic SSRI Trial I: D	methorphan and bupropion inadequate response at a than isomer, prodrug or metal than isomer, prodrug or metal drug of the same chemical quired trials may be overrided lically contraindicated. Fetzima Other: Tructions Trug Name & Dose Trug Name & Dose	extended- rele erapeutic dose bolite of the re I entity that res den when docur Quantity	ase tablet (with an ext quested me sulted in a p mented evid	Auvelity), one of the ended-release buping dication, one of the edication, one of the edication artial response with dence is provided the ended the edication are provided the edication are provided the edication are provided the edication are edicated as a constant of the edicated	e trials ropion e trials h a
must include a previous trial and agent; and 9) If the request is for must be with the preferred parent documented intolerance. The requise of these agents would be med Non-Preferred Aplenzin Auvelity Strength Dosage Instruction Diagnosis: Preferred Generic SSRI Trial I: Defailure Reason Preferred Generic SSRI Trial 2: Defailure Reason Preferred Generic SNRI Trial 2: Defailure Reason Preferred Generic SNRI Trial 2: Defailure Reason	methorphan and bupropion inadequate response at a than isomer, prodrug or metal to drug of the same chemical quired trials may be override lically contraindicated. Fetzima Other:	extended- rele erapeutic dose bolite of the re I entity that res den when docur Quantity	ase tablet (with an ext quested me sulted in a p mented evid Days Su	Auvelity), one of the ended-release bupinedication, one of the edication, one of the eartial response with dence is provided the experience of the ended of the end of th	e trials ropion e trials h a
must include a previous trial and agent; and 9) If the request is for must be with the preferred parent documented intolerance. The requese of these agents would be med Non-Preferred Aplenzin Auvelity Strength Dosage Instruction Diagnosis: Preferred Generic SSRI Trial I: Defailure Reason Preferred Generic SSRI Trial 2: Defailure Reason Preferred Generic SNRI Trial: Defailure Reason Preferred Generic SNRI Trial: Defailure Reason	methorphan and bupropion inadequate response at a than isomer, prodrug or metal at drug of the same chemical quired trials may be overrided lically contraindicated. Fetzima Other:	extended- rele erapeutic dose bolite of the re I entity that res den when docur Quantity	ase tablet (with an ext quested me sulted in a p mented evid	Auvelity), one of the ended-release bupinedication, one of the edication, one of the edication artial response with dence is provided the ended of the edication are provided the edication are edicated as a constant of the edicate edicated are edicated as a constant of the edicated a	e trials ropion e trials h a
must include a previous trial and agent; and 9) If the request is for must be with the preferred parent documented intolerance. The requise of these agents would be med Non-Preferred Aplenzin Auvelity Strength Dosage Instruction Diagnosis: Preferred Generic SSRI Trial I: Defailure Reason Preferred Generic SSRI Trial 2: Defailure Reason Preferred Generic SNRI Trial 2: Defailure Reason Preferred Generic SNRI Trial 2: Defailure Reason	methorphan and bupropion inadequate response at a than isomer, prodrug or metal than isomer, prodrug be override lically contraindicated. Fetzima	extended- rele erapeutic dose bolite of the re I entity that res den when docur Quantity Quantity	ase tablet (with an ext quested me sulted in a p mented evid Days Su	Auvelity), one of the ended-release bupiedication, one of the edication, one of the edication artial response with dence is provided the ended the edication artial Dates: Trial Dates: Trial Dates: Trial Dates:	e trials ropion e trials h a
must include a previous trial and agent; and 9) If the request is for must be with the preferred parendocumented intolerance. The requese of these agents would be med Non-Preferred Aplenzin Auvelity Strength Dosage Instruction Diagnosis: Preferred Generic SSRI Trial I: Defailure Reason Preferred Generic SSRI Trial 2: Defailure Reason Preferred Generic SNRI Trial: Defailure Reason	methorphan and bupropion inadequate response at a than isomer, prodrug or metal at drug of the same chemical quired trials may be override lically contraindicated. Fetzima Other: ructions rug Name & Dose rug Name & Dose grug Name & Dose drug Name & Dose	extended- rele erapeutic dose bolite of the re I entity that res den when docur Quantity Quantity Solution of the release o	ase tablet (with an ext quested me sulted in a p mented evid Days Su Tr	Auvelity), one of the ended-release bupinedication, one of the edication, one of the edication artial response with dence is provided the ended of the edication are provided the edication artial Dates: Trial Dates:	e trials ropion e trials h a



Request for Prior Authorization

FAX Completed Form To 1 (800) 574-2515

> Provider Help Desk | (877) 776-1567

ANTIDEPRESSANTS (PLEASE PRINT – ACCURACY IS IMPORTANT)

Vortioxetine Trial: Dose	Trial Dates:	
Failure Reason		
Antidepressant plus adjunct trials:		
Antidepressant Trial: Drug Name & Dose	Trial Dates:	
Failure Reason		
Adjunct Trial: Drug Name & Dose	Trial Dates:	
Failure Reason		
Requests for Auvelity:		
Extended-Release Bupropion Trial: Dose	Trial Dates:	
Failure Reason		
Medical or contraindication reason to override trial requirements:		
Attach lab results and other documentation as necessary.		
Prescriber signature (Must match prescriber listed above.)	Date of submission	

IMPORTANT NOTE: In evaluating requests for prior authorization the consultant will consider the treatment from the standpoint of medical necessity only. If approval of this request is granted, this does not indicate that the member continues to be eligible for Medicaid. It is the responsibility of the provider who initiates the request for prior authorization to establish by inspection of the member's Medicaid eligibility card and, if necessary by contact with the county Department of Health and Human Services, that the member continues to be eligible for Medicaid.