

Iowa Department of Human Services

Request for Prior Authorization Finerenone (Kerendia)



(PLEASE PRINT - ACCURACY IS IMPORTANT)

FAX Completed Form To 1 (877) 733-3195 Provider Help Desk 1 (844) 236-1464

IA Medicaid Member ID #	Patient name	DOB		
Patient address				
Provider NPI	Prescriber name	Phone		
Prescriber address		Fax		
Pharmacy name	Address	Phone		
Prescriber must complete all information above. It must be legible, correct, and complete or form will be returned.				
Pharmacy NPI	Pharmacy fax ND0	° │		

Prior authorization (PA) is required for finerenone (Kerendia). Payment will be considered under the following conditions:

- 1) Request adheres to all FDA approved labeling for indication, including age, dosing, contraindications, warnings and precautions, and drug interactions; and
- 2) Patient has a diagnosis of chronic kidney disease (CKD) associated with Type 2 Diabetes (T2D); and
- 3) Patient is currently receiving a maximally tolerated dose of an angiotensin converting enzyme inhibitor (ACEi) or angiotensin receptor blocker (ARB); and
- 4) Patient is currently receiving a maximally tolerated dose of a sodium-glucose co-transporter 2 (SGLT2) inhibitor indicated to reduce the risk of sustained eGFR decline, end-stage kidney disease, cardiovascular death, and hospitalization for heart failure in adults with chronic kidney disease [i.e., dapagliflozin (Farxiga)]; and
- 5) Patient has the following baseline tests prior to initiation of treatment with finerenone:
 - a. Serum potassium is ≤ 5.0 mEq/L; and
 - b. Estimated glomerular filtration rate (eGFR) is \geq 25 mL/min/1.73m²; and
 - c. Urine albumin to creatinine ration (UACR) is \geq 30 mg/g.

The required trials may be overridden when documented evidence if provided that the use of these agents would be medically contraindicated.

Initial authorizations will be approved for six months. Additional PAs will be considered with the following documentation:

- 1. Patient's serum potassium is < 5.5 mEq/L; and
- 2. Patient's eGFR is \geq 25 mL/min/1.73m2; and
- 3. Patient remains on a maximally tolerated dose of an ACEi or ARB; and
- 4. Patient remains on a maximally tolerated dose of an SGLT2 inhibitor.

Non-Preferred

Kerendia

	Strength	Dosage Instructions	Quantity	Days Supply
anosis:	<u> </u>			

lov	va Departmer	nt of Human Se	ervices	
Request		uthorization ne (Kerendia		
(PLEA		•	S IMPORTANT)	
Document current treatment of a maxim	ally talarata	daga of an A		
Drug Name & Dose:	•			
Document current treatment of a maxim sustained eGFR decline, end-stage kidn adults with chronic kidney disease: Drug Name & Dose:	ey disease, o	cardiovascula	r death, and hospitalizatio	on for heart failure in
Baseline tests prior to initiation of treatr	nent (attach	results):		
	Yes			
	Yes	🛛 No		
 UACR ≥ 30mg/g 	Yes	🛛 No		
Renewal Requests				
Updated tests (attach results)				
○ Serum Potassium < 5.5 mEq/L	Yes	🛛 No		
 eGFR ≥ 25mL/min/1.73m₂ 	Yes	🛛 No		
Patient remains on a maximally tolerated	d dose of AC	Ei or ARB:		
Yes Drug Name & Dose:				
D No				
Patient remains on a maximally tolerated eGFR decline, end-stage kidney disease chronic kidney disease:				
Yes Drug Name & Dose:				
□ No				

Attach lab results and other documentation as necessary.

Prescriber signature (Must match prescriber listed above.)	Date of submission

IMPORTANT NOTE: In evaluating requests for prior authorization the consultant will consider the treatment from the standpoint of medical necessity only. If approval of this request is granted, this does not indicate that the member continues to be eligible for Medicaid. It is the responsibility of the provider who initiates the request for prior authorization to establish by inspection of the member's Medicaid eligibility card and, if necessary by contact with the county Department of Human Services, that the member continues to be eligible for Medicaid.