



Original Effective Date: 06/2013  
Current Effective Date: 01/07/2023  
Last P&T Approval/Version: 10/26/2022  
Next Review Due By: 10/2023  
Policy Number: C2729- A

## Krystexxa (pegloticase)

### PRODUCTS AFFECTED

Krystexxa (pegloticase)

### COVERAGE POLICY

Coverage for services, procedures, medical devices and drugs are dependent upon benefit eligibility as outlined in the member's specific benefit plan. This Coverage Guideline must be read in its entirety to determine coverage eligibility, if any.

This Coverage Guideline provides information related to coverage determinations only and does not imply that a service or treatment is clinically appropriate or inappropriate. The provider and the member are responsible for all decisions regarding the appropriateness of care. Providers should provide Molina Healthcare complete medical rationale when requesting any exceptions to these guidelines.

#### **Documentation Requirements:**

Molina Healthcare reserves the right to require that additional documentation be made available as part of its coverage determination; quality improvement; and fraud; waste and abuse prevention processes. Documentation required may include, but is not limited to, patient records, test results and credentials of the provider ordering or performing a drug or service. Molina Healthcare may deny reimbursement or take additional appropriate action if the documentation provided does not support the initial determination that the drugs or services were medically necessary, not investigational or experimental, and otherwise within the scope of benefits afforded to the member, and/or the documentation demonstrates a pattern of billing or other practice that is inappropriate or excessive.

#### **DIAGNOSIS:**

Chronic Gout

#### **REQUIRED MEDICAL INFORMATION:**

This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. If a drug within this policy receives an updated FDA label within the last 180 days, medical necessity for the member will be reviewed using the updated FDA label information along with state and federal requirements, benefit being administered and formulary preferencing. Coverage will be determined on a case-by case basis until the criteria can be updated through Molina Healthcare, Inc. clinical governance. Additional information may be required on a case-by-case basis to allow for adequate review.

#### **A. CHRONIC GOUT**

1. Documented diagnosis of gout  
AND
2. Documentation of ONE of the following:
  - (a) at least 2 gout flares in the previous 12 months  
OR
  - (b) a history of at least 1 gouty tophus  
OR

Molina Healthcare, Inc. confidential and proprietary © 2022

This document contains confidential and proprietary information of Molina Healthcare and cannot be reproduced, distributed, or printed without written permission from Molina Healthcare. This page contains prescription brand name drugs that are trademarks or registered trademarks of pharmaceutical manufacturers that are not affiliated with Molina Healthcare.

## Drug and Biologic Coverage Criteria

(c) chronic gouty arthropathy

AND

3. Documentation of baseline serum uric acid level >6mg/dL [DOCUMENTATION REQUIRED]  
AND
4. Documentation of a therapeutic trial and clinical failure of TWO of the following for maintenance treatment of gout, at a maximum tolerated dose or an adequate therapeutic dose for 3 months: allopurinol [Maximum Doses: Alopurinol® 600 mg/day; Zylor® 800 mg/day] OR a probenecid containing medication (e.g., probenecid or probenecid-colchicine)] OR febuxostat  
*\*\*\*Clinical failure is defined as the inability to maintain serum uric acid (SUA) level  $\leq 6$  mg/dL after reaching the maximum tolerated dose or an adequate therapeutic dose\*\*\**  
AND
5. Prescriber attests that member will NOT concurrently be receiving other urate lowering therapies such as allopurinol, febuxostat, probenecid, lesinurad, etc.  
AND
6. Documentation that member has been screened for higher risk for glucose 6 phosphate dehydrogenase (G6PD) deficiency and found negative for G6PD before starting Krystexxa.  
AND
7. Prescriber attests that member will be pre-medicated with antihistamines and corticosteroids and closely monitored for anaphylaxis for an appropriate period of time after administration of Krystexxa. Note: Anaphylaxis may occur with any infusion, including a first infusion, and generally manifests within 2 hours of the infusion. However, delayed type hypersensitivity reactions have also been reported.

## CONTINUATION OF THERAPY:

### A. CHRONIC GOUT

1. Adherence to therapy at least 85% of the time as verified by the prescriber or member medication fill history OR adherence less than 85% of the time due to the need for surgery or treatment of an infection, causing temporary discontinuation  
AND
2. Prescriber attests to or clinical reviewer has found no evidence of intolerable adverse effects or drug toxicity  
AND
3. Documented positive response to Krystexxa® (pegloticase) treatment, including, but not limited to: reduction in serum uric acid levels compared to baseline or reductions in gout flares  
[DOCUMENTATION REQUIRED]  
AND
4. Prescriber attests that the member has not had 2 consecutive uric acid levels >6 mg/dL while on therapy  
NOTE: Members with documented serum uric acid concentrations (prior to their next infusion) of greater than 6 mg/dl on more than one occasion during treatment must discontinue treatment with pegloticase.  
The risk of infusion reaction is higher in members whose uric acid level increases to above 6 mg/dL, particularly when 2 consecutive levels above 6 mg/dL are observed. Monitor serum uric acid levels prior to infusions and consider discontinuing treatment if levels increase to above 6 mg/dL despite treatment.  
AND
5. Prescriber attests that member will be pre-medicated with antihistamines and corticosteroids and closely monitored for anaphylaxis for an appropriate period of time after administration of Krystexxa. Note: Anaphylaxis may occur with any infusion, and generally manifests within 2 hours of the infusion. However, delayed type hypersensitivity reactions have also been reported.

## DURATION OF APPROVAL:

Initial authorization: 6 months, Continuation of therapy: 12 months

## Drug and Biologic Coverage Criteria

### **PRESCRIBER REQUIREMENTS:**

Prescribed by or in consultation with a rheumatologist or nephrologist [If prescribed in consultation, consultation notes must be submitted with initial request and reauthorization requests]

### **AGE RESTRICTIONS:**

18 years of age or older

### **QUANTITY:**

Quantity limitation: maximum of twelve (12) infusions for 6 months; maximum of twenty-six (26) infusions for 12 months. [Krystexxa 8mg IV every two (2) weeks. Dose not to exceed one 8mg infusion every 2 weeks]

### **PLACE OF ADMINISTRATION:**

The recommendation is that infused medications in this policy will be for pharmacy or medical benefit coverage administered in a place of service that is a non-hospital facility-based location as per the Molina Health Care Site of Care program.

**Note:** Site of Care Utilization Management Policy applies for Krystexxa (pegloticase). For information on site of care, see

[Specialty Medication Administration Site of Care Coverage Criteria \(molinamarketplace.com\)](https://molinahealthcare.com/specialty-medication-administration-site-of-care-coverage-criteria)

## **DRUG INFORMATION**

### **ROUTE OF ADMINISTRATION:**

Intravenous

### **DRUG CLASS:**

Gout Agents

### **FDA-APPROVED USES:**

For the treatment of chronic gout in adult patients refractory to conventional therapy.

### **COMPENDIAL APPROVED OFF-LABELED USES:**

None

## **APPENDIX**

### **APPENDIX:**

None

## **BACKGROUND AND OTHER CONSIDERATIONS**

### **BACKGROUND:**

Pegloticase (Krystexxa™) has been indicated for the treatment of chronic gout in adult members refractory to conventional therapy. Pegloticase (Krystexxa™) is a PEGylated uric acid-specific enzyme that reduces serum uric acid levels by catalyzing the oxidation of uric acid to allantoin. Pegloticase is a PEGylated uric acid-specific enzyme that consists of recombinant modified mammalian urate oxidase produced by a genetically modified strain of Escherichia coli (Krystexxa prescribing information, 2010). It is approved for the treatment of chronic gout in adult member s' refractory to conventional therapy.

Krystexxa is not recommended for the treatment of asymptomatic hyperuricemia

### **CONTRAINDICATIONS/EXCLUSIONS/DISCONTINUATION:**

All other of Krystexxa (pegloticase) are considered experimental/investigational and therefore, will follow Molina's Off- Label policy

## Drug and Biologic Coverage Criteria

Contraindications to (Krystexxa (pegloticase) include: Patients with glucose-6-phosphate dehydrogenase (G6PD) deficiency, patients with history of serious hypersensitivity reactions, including anaphylaxis, to KRYSTEXXA or any of its components

### OTHER SPECIAL CONSIDERATIONS:

Krystexxa (pegloticase) has a black box warning for anaphylaxis and infusion reactions, G6PD deficiency associated hemolysis and methemoglobinemia.

### CODING/BILLING INFORMATION

*Note: 1) This list of codes may not be all-inclusive. 2) Deleted codes and codes which are not effective at the time the service is rendered may not be eligible for reimbursement*

HCPSC CODE	DESCRIPTION
J2507	Injection, pegloticase, 1 mg

### AVAILABLE DOSAGE FORMS:

Krystexxa SOLN 8MG/ML (1ml vial)

### REFERENCES

1. Krystexxa [package insert]. Deerfield, IL: Horizon Therapeutics USA Inc; July 2022.
2. Khanna, D., et al. 2012 American College of Rheumatology Guideline Management of Gout part 1. Arthritis Care & Research: Vol 64, No 10, October 2012, pp 1431-1446.
3. Sivera F, Andres M, Carmona L et al. Recommendations for the Diagnosis and Management of Gout. Ann Rheum Dis 2014; 73(2):328-335.
4. Qaseem Amir, et al. Management of Acute and Recurrent Gout: A Clinical Practice Guideline from the American College of Physicians. Ann Intern Med. Doi: 10.7326/M16-0570. November 2016.
5. FitzGerald, J., Dalbeth, N., Mikuls, T., Brignardello-Petersen, R., Guyatt, G., & Abeles, A. et al. (2020). 2020 American College of Rheumatology Guideline for the Management of Gout. Arthritis Care & Research, 72(6), 744-760. doi: 10.1002/acr.24180

SUMMARY OF REVIEW/REVISIONS	DATE
REVISION- Notable revisions: Required Medical Information Continuation of Therapy Prescriber Requirements FDA-Approved Uses Contraindications/Exclusions/Discontinuation Other Special Considerations References	Q4 2022
Q2 2022 Established tracking in new format	Historical changes on file