

Cardio Policy:

Renal/Retroperitoneal Vascular Duplex Ultrasound

POLICY NUMBER UM CARDIO_1125	SUBJECT Renal/Retroperitoneal Vascular Duplex Ultrasound		DEPT/PROGRAM UM Dept	PAGE 1 OF 4
DATES COMMITTEE REVIEWED 07/22/11, 12/12/12, 06/30/14, 08/22/14, 08/12/15, 11/28/16, 12/21/16, 10/10/17, 02/13/19, 03/08/19, 09/11/19, 12/11/19, 08/12/20, 04/14/21, 08/11/21, 09/08/21, 12/08/21, 12/14/22, 05/10/23, 12/20/23	APPROVAL DATE December 20, 2023	EFFECTIVE DATE December 22, 2023	COMMITTEE APPROVAL DATES 07/22/11, 12/12/12, 06/30/14, 08/22/14, 08/12/15, 11/28/16, 12/21/16, 10/10/17, 02/13/19, 03/08/19, 09/11/19, 12/11/19, 08/12/20, 04/14/21, 08/11/21, 09/08/21, 12/08/21, 12/14/22, 05/10/23, 12/20/23	
PRIMARY BUSINESS OWNER: UM		COMMITTEE/BOARD APPROVAL Utilization Management Committee		
URAC STANDARDS HUM v8: UM 1-2; UM 2-1	NCQA STANDARDS UM 2		ADDITIONAL AREAS OF IMPACT	
CMS REQUIREMENTS	STATE/FEDERAL REQUIREMENTS		APPLICABLE LINES OF BUSINESS Commercial, Exchange, Medicaid	

I. PURPOSE

Indications for determining medical necessity for Renal/Retroperitoneal Vascular Duplex Ultrasound.

II. DEFINITIONS

Renal Duplex ultrasound images the renal arteries via spectral/color flow Doppler and B-mode scanning to assess abnormalities in the blood flow there by identifying areas of blockage.

An appropriate diagnostic or therapeutic procedure is one in which the expected clinical benefit exceeds the risks or negative consequences of the procedure by a sufficiently wide margin such that the procedure is generally considered acceptable or reasonable care. The ultimate objective of AUC is to improve patient care and health outcomes in a cost–effective manner but is not intended to ignore ambiguity and nuance intrinsic to clinical decision making.

Appropriate Care - Median Score 7-9

May be Appropriate Care - Median Score 4-6

Rarely Appropriate Care - Median Score 1-3

III. POLICY

Indications for approving a request for medical necessity:

A. Follow up testing in asymptomatic patient with Renal Artery Stenosis to determine hemodynamic significance of stenosis. (AUC Score 7)^{1,2,3,4,5,6,8}

- B. Evaluation for Renal Artery Stenosis with worsening of renal function test (Creatinine) in presence of malignant hypertension. (AUC Score 8)1,2,3,4,5,6,8
- C. Evaluation for Renal Artery Stenosis with resistant hypertension (patient has uncontrolled hypertension after being on at least 3 medications including diuretics). (AUC Score 8)^{1,2,3,4,5,6,8}
- D. Evaluation for Renal Artery Stenosis in an asymptomatic patient with hypertension who in younger than 30 years old. (AUC Score 8)1,2,3,4,5,6,8
- E. Evaluation for Renal Artery Stenosis in symptomatic patient, with elevated creatinine and hypertension in young person (age less than 35 years). (AUC Score 8)^{1,2,3,4,5,6,8}
- F. Evaluation for Renal Artery Stenosis in symptomatic patient, with elevated creatinine and hypertension with evidence of size discrepancy between kidneys noted on other imaging modalities as CT/MR. (AUC Score 7)1,2,3,4,5,8
- G. Evaluation for Renal Artery Stenosis in symptomatic patient, with elevated creatinine and hypertension. (AUC Score 7)^{1,2,3,4,5,8}
- H. Evaluation for Renal Artery Stenosis in symptomatic patient, with elevated creatinine (greater than 50% increase from baseline) and/or hypertension with recent addition of ACEI/ARBs. (AUC Score 8)1,2,3,4,5,8
- I. Evaluation for Renal Artery Stenosis in symptomatic patient, with elevated creatinine and/or hypertension with epigastric bruit. (AUC Score 7)1,2,3,4,5,8
- J. Evaluation for Renal Artery Stenosis in symptomatic patient, with elevated creatinine and/or hypertension (AUC Score 8)1,2,3,4,5,8
- K. Baseline surveillance within one month after Renal Artery revascularization. (AUC Score 8)1,2,3,4,5,8
- L. Renal Duplex in a patient after Renal Artery Revascularization with new or worsening symptoms related to renal artery stenosis. (AUC Score 8)^{1,2,3,4,5,8}
- M. To evaluate patients presenting with signs or symptoms such as epigastric or periumbilical postprandial pains that last for 1-3 hours and/or with associated weight loss resulting from decreased oral intake which may indicate chronic intestinal (mesenteric or celiac artery) ischemia. (AUC Score 6)^{1,2,3,4,5,8}
- N. Baseline surveillance in asymptomatic patients one month after mesenteric revascularization, and once again at 3-5 months post, 6-8 months post, and 9-12 months post (AUC Score 8)^{1,2,3,4,5,8} imaging beyond one year is rarely necessary.
- O. To evaluate for suspected portal hypertension or portal vein thrombosis in the presence of hepatic disease, no prior ultrasound within the last 6 months. (AUC Score 7)^{1,2,3,4,5,8}
- P. Surveillance ultrasound to assess for complications status post hepatic or renal or pancreas transplant may be performed as per the protocol of the transplant facility. (AUC Score 7)^{7,8}
- Q. Surveillance ultrasound to assess for development of hepatocellular carcinoma may be performed every 6 months in patients with the following primary liver conditions: Hepatitis C, EtOH liver disease, hereditary hemochromatosis, primary biliary cholangitis, autoimmune hepatitis, or alpha 1-atitrypsin deficiency. (AUC Score 6)⁹
- R. To evaluate patients with pain or swelling of scrotal contents which may be as a result of suspected obstruction in arterial inflow or venous outflow to the testicles or related structures. The

- use of duplex scanning of scrotal contents should only be performed after conventional diagnostic test, such as ultrasound, have proven to be "non-definitive." (AUC Score 7)¹⁰
- S. Renal Duplex is appropriate to perform, to evaluate retroperitoneal vasculature structures as initial workup, prior to any organ transplant, no prior ultrasound within the last 6 months. (AUC Score 7) 12,13,14,15

Limitations:

- A. Screening of asymptomatic patients for Renal Artery Stenosis with evidence of atherosclerotic vascular disease in other beds is inappropriate.
- B. Screening of asymptomatic patients for Renal Artery Stenosis with unexplained size discrepancy between kidneys noted on other imaging modalities is inappropriate.
- C. Follow up testing for Renal Artery Stenosis as surveillance in asymptomatic patients is inappropriate.
- D. Requests for services that are part of a surveillance protocol for patients who are involved in a clinical trial are considered out of scope (OOS) for New Century Health and cannot be reviewed.

IV. PROCEDURE

- A. In order to review a request for medical necessity, the following items must be submitted for review:
 - 1. Progress note that prompted request
 - 2. Most recent renal duplex ultrasound of CT/MR report (if applicable)
- B. Primary code appropriate for this service:93975 for complete study of the abdominal, pelvic, scrotal contents and/or retroperitoneal organs93976 for limited study.

V. APPROVAL AUTHORITY

- A. Review Utilization Management Department
- B. Final Approval Utilization Management Committee

VI. ATTACHMENTS

A. None

VII. REFERENCES

- 1. Centers for Medicare and Medicaid Services. Florida Local Coverage Determination (LCD) (L33674). Duplex Scanning. Retrieved from https://www.cms.gov [Accessed December 19, 2023].
- 2. Centers for Medicare and Medicaid Services. Michigan Local Coverage Determination (LCD) (L35755). Duplex Scanning. Retrieved from https://www.cms.gov [Accessed December 19, 2023].
- Centers for Medicare and Medicaid Services. Illinois Local Coverage Determination (LCD) (L33627). Duplex Scanning. Retrieved from https://www.cms.gov [Accessed December 19, 2023].
- Anderson, JL, et al. Management of Patients with Peripheral Artery Disease (Compilation of 2005 and 2011 ACCF/AHA Guideline Recommendations) A Report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines

- Developed in Collaboration with the Society for Cardiovascular Angiography and Interventions, Society of Interventional Radiology, Society for Vascular Medicine, and Society for Vascular Surgery. JACC Vol. 61, No. 14, 2013 pp 1555-1570
- 5. Emile R. Mohler, III, MD, FACC, et. al. ACCF/ACR/AIUM/ASE/ASN/ICAVL/SCAI/SCCT/SIR/SVM/SVS 2012 appropriate use criteria for peripheral vascular ultrasound and physiological testing part I: Arterial ultrasound and physiological testing. A Report of the American College of Cardiology Foundation Appropriate Use Criteria Task Force, American College of Radiology, American Institute of Ultrasound in Medicine, American Society of Echocardiography, American Society of Nephrology, Intersocietal Commission for the Accreditation of Vascular Laboratories, Society for Cardiovascular Angiography and Interventions, Society of Cardiovascular Computed Tomography, Society for Interventional Radiology, Society for Vascular Medicine, and Society for Vascular Surgery. Journal of the American College of Cardiology, July 2012, Volume 60, Issue 3, Pages 242-276.
- 6. Simonetto, DA et al. ACG Clinical Guideline: Disorders of the Hepatic and Mesenteric Circulation. The Amer Journal of Gastro. January 2020 Volume 115 Issue 1 pp 18-40
- 7. United Network for Organ Sharing (UNOS). United Network for Organ Sharing (UNOS). Data. UNOS Web site. Available at http://Available at https://unos.org/data/. March 20, 2018; Accessed: : March 1, 2019.
- 8. Robert C. Hendel MD, FACC, FAHA, et al. Appropriate use of cardiovascular technology: 2013 ACCF appropriate use criteria methodology update: a report of the American College of Cardiology Foundation appropriate use criteria task force. Journal of the American College of Cardiology, March 2013. Volume 61. Issue 12. Pages 1305-1317.
- 9. Patrick S Harris, et al. Hepatocellular carcinoma surveillance: An evidence-based approach. World J Gastroenterol. 2019 Apr 7; 25(13): 1550–1559.
- 10. SizeWu, et al. Curated findings and implications in duplex ultrasound interrogation of the scrotum or varicoceles. Scientific Reports | (2020) 10:22028 | https://doi.org/10.1038/s41598-020-78619-1
- 11. NCQA UM 2023 Standards and Elements.
- Pre-Renal Transplant Risk Stratification. J Am Coll Cardiol Img. 2018 Jun, 11 (6) 855–858.https://doi.org/10.1016/j.jcmg.2017.08.001
- 13. DeBolle SA, Ochieng IA, Saha AK, Sung RS. Evaluation of the Effectiveness of Screening for Iliac Arterial Calcification in Kidney Transplant Candidates. Ann Transplant. 2020 Sep 15;25:e922178. doi: 10.12659/AOT.922178. PMID: 32929057; PMCID: PMC7518019.
- Chadban, Steven J. BMed, PhD1Gregory A. MD, MSc15,*. KDIGO Clinical Practice Guideline on the Evaluation and Management of Candidates for Kidney Transplantation. Transplantation: April 2020 - Volume 104 - Issue 4S1 - p S11-S103
- Martin, P., DiMartini, A., Feng, S., Brown, R., Jr. and Fallon, M. (2014), Evaluation for liver transplantation in adults: 2013 practice guideline by the American Association for the Study of Liver Diseases and the American Society of Transplantation. Hepatology, 59: 1144-1165. https://doi.org/10.1002/hep.26972