

HEDIS® Tips:

Chlamydia Screening in Women (CHL)

MEASURE DESCRIPTION

The percentage of women 16-24 years of age who were identified as sexually active and who had at least one chlamydia test during the measurement year.

Exclusion: Patients who were included in the measure based on pregnancy test alone and on the date of the pregnancy test or the 6 days after the pregnancy test, had a prescription for isotretinoin or an x-ray.

CODES INCLUDED IN THE CURRENT HEDIS MEASURES

Codes to Identify Chlamydia Screening

Description	CPT® Codes
Chlamydia Tests	87110, 87270, 87320, 87490 87492, 87810

HOW TO IMPROVE HEDIS SCORES

- ☐ Perform chlamydia screening every year on every 16-24 year old female identified as sexually active (use any visit opportunity).
- ☐ Add chlamydia screening as a standard lab for women 16-24 years old. Use well child exams and well women exams for this purpose.
- ☐ Ensure that you have an opportunity to speak with your adolescent female patients without her parent.
- ☐ Remember that chlamydia screening can be performed through a urine test. Offer this as an option for your patients.
- ☐ Place chlamydia swab next to Pap test or pregnancy detection materials.

The information presented herein is for informational and illustrative purposes only. It is not intended, nor is it to be used, to define a standard of care or otherwise substitute for informed medical evaluation, diagnosis and treatment which can be performed by a qualified medical professional. Molina Healthcare Inc. does not warrant or represent that the information contained herein is accurate or free from defects.

All summaries of the measures contained herein are reproduced with permission from HEDIS Volume 2: Technical Specifications for Health Plans by the National Committee for Quality Assurance (NCQA). HEDIS® is a registered trademark of the National Committee for Quality Assurance (NCQA). Updated 01/16/2020