

# HEDIS® Tips:

## Immunizations for Adolescents (IMA)

### MEASURE DESCRIPTION

Adolescents 13 years of age who received the following vaccines **on or before the 13<sup>th</sup> birthday**:

- One meningococcal conjugate vaccine (must be completed on or between the 11<sup>th</sup> and 13<sup>th</sup> birthdays)
- One tetanus, diphtheria toxoids and acellular pertussis (Tdap) (must be completed on or between the 10<sup>th</sup> and 13<sup>th</sup> birthdays)
- At least two Human Papillomavirus (HPV) vaccines or three HPV vaccines with different dates of service on or between the 9<sup>th</sup> and 13<sup>th</sup> birthdays

### USING CORRECT BILLING CODES

#### Codes to Identify Adolescent Immunizations

Description	CPT® /CVX Codes
Meningococcal	<b>CPT:</b> 90734 <b>CVX:</b> 108,114,136, 147, 167
Tdap	<b>CPT:</b> 90715 <b>CVX:</b> 115
HPV	<b>CPT:</b> 90649,90650, 90651 <b>CVX:</b> 62,118, 137,165

### HOW TO IMPROVE HEDIS SCORES

- ☐ Use the State immunization registry.
- ☐ Review missing vaccines with parents.
- ☐ Recommend immunizations to parents. Parents are more likely to agree with vaccinations when supported by the provider. Address common misconceptions about vaccinations.
- ☐ Train office staff to prep the chart in advance of the visit and identify overdue immunizations.
- ☐ Make every office visit count- take advantage of sick visits for catching up on needed vaccines.
- ☐ Institute a system for patient reminders.
- ☐ Some vaccines may have been given before patients were Molina members. Include these on the members' vaccination record even if your office did not provide the vaccine.
- ☐ Administer the HPV vaccine at the same time as other vaccines. Inform parents that the full HPV vaccine series requires 2 or 3 shots and have a system for patient reminders.
- ☐ Recommend the HPV vaccine series the same way you recommend other adolescent vaccines. Parents are more likely to agree with vaccinations when supported by the provider. Address common misconceptions about the HPV vaccine.