HEDIS[®] Tips:

Statin Therapy for Patients with Cardiovascular Disease (SPC)

MEASURE DESCRIPTION

The percentage of males 21-75 years and females 40-75 years who were identified as having clinical atherosclerotic cardiovascular disease (ASCVD) and met the following criteria:

- Received Statin Therapy. Patients were dispensed at least one high or moderate-intensity statin medication during the measurement year.
- Statin Adherence 80%. Patients remained on a high or moderate-intensity statin medication for at least 80% of the treatment period.

USING CORRECT PRESCRIPTIONS

High and Moderate-Intensity Statin Medications

Description	Prescription	
High-intensity statin therapy	Atorvastatin 40–80 mg	Rosuvastatin 20–40 mg
	Amlodipine-atorvastatin 40-80 mg	Simvastatin 80 mg
	Ezetimibe-atorvastatin 40-80 mg	• Ezetimibe-simvastatin 80 mg
Moderate-intensity statin therapy	Atorvastatin 10–20 mg	Sitagliptin-simvastatin 20-40 mg
	Amlodipine-atorvastatin 10-20 mg	Pravastatin 40–80 mg
	• Ezetimibe-atorvastatin 10-20 mg	Lovastatin 40 mg
	Rosuvastatin 5–10 mg	Niacin-lovastatin 40 mg
	Simvastatin 20–40 mg	Fluvastatin XL 80 mg
	• Ezetimibe-simvastatin 20-40 mg	Fluvastatin 40 mg bid
	Niacin-simvastatin 20-40 mg	Pitavastatin 2–4 mg

*Please refer to the Molina Healthcare Drug Formulary at <u>www.molinahealthcare.com</u>for statin medications that may require prior authorization or step therapy.

HOW TO IMPROVE HEDIS SCORES

Continue to stress the value of prescribed medications for managing cardiovascular disease and the importance of adherence throughout the entire treatment period.

Schedule appropriate follow-up with patients to assess if medication is taken as prescribed.

- Do not rely on the patient to follow through with scheduling subsequent appointments. Routinely arrange the next appointment when the patient is in the office. If the patient misses a scheduled appointment, office staff should contact the patient to assess why appointment was missed.
- □ Contact Health Care Services at your affiliated Molina Healthcare State plan for additional information about Medication Therapy Management (MTM) criteria and to request a referral for patients with at least six (6) chronic medications and at least three (3) qualifying diagnoses. They may be eligible for MTM sessions.
- Provide smoking cessation and other interventions to eliminate or control risk factors.



