

HEDIS® Tips:

Initiation & Engagement of Alcohol & Other Drug Abuse or Dependence Treatment (IET)

MEASURE DESCRIPTION

The percentage of adolescent and adult patients 13 years of age and older with a new episode of alcohol or other drug (AOD) abuse or dependence with the following:

- **Initiation of AOD Treatment.** Initiate treatment through inpatient AOD admission, outpatient visit, intensive outpatient encounter or partial hospitalization, telehealth, or medication treatment within 14 days of diagnosis.
- **Engagement of AOD Treatment.** Initiated treatment and had two or more additional AOD services or medication treatment within 34 days of the initiation visit.

USING CORRECT BILLING CODES

Codes to Identify AOD Abuse or Dependence

ICD-10-CM Diagnosis

F10.10 F10.120, F10.121, F10.129, F10.14, F10.150, F10.151, F10.159, F10.180-F10.182, F10.188, F10.19-F10.20, F10.220, F10.221, F10.229-F10.232, F10.239, F10.24, F10.250-F10.251, F10.259, F10.26, F10.27, F10.280-F10.282, F10.288, F10.29, F11.10, F11.120-F11.122, F11.129, F11.14, F11.150, F11.151, F11.159, F11.181, F11.182, F11.188, F11.19, F11.20, F11.220-F11.222, F11.229, F11.23, F11.24, F11.250, F11.251, F11.259, F11.281, F11.282, F11.288, F11.29, F12.10, F12.120-F12.122, F12.129, F12.150, F12.151, F12.159, F12.180, F12.188, F12.19, F12.20, F12.220-F12.222, F12.229, F12.23, F12.250, F12.251, F12.259, F12.280, F12.288, F12.29, F13.10, F13.120, F13.121, F13.129, F13.14, F13.150, F13.151, F13.159, F13.180- F13.182, F13.188, F13.19, F13.20, F13.220, F13.221, F13.229-F13.232, F13.239, F13.24, F13.250, F13.251, F13.259, F13.26, F13.27, F13.280-F13.282, F13.288, F13.29, F14.10, F14.120-F14.122, F14.129, F14.14, F14.150, F14.151, F14.159, F14.180-F14.182, F14.188, F14.19, F14.20, F14.220-F14.222, F14.229, F14.23, F14.24, F14.250, F14.251, F14.259, F14.280-F14.282, F14.288, F14.29, F15.10, F15.120-F15.122, F15.129, F15.14, F15.150, F15.151, F15.159, F15.180- F15.182, F15.188, F15.19, F15.20, F15.220-F15.222, F15.229, F15.23, F15.24, F15.250, F15.251, F15.259, F15.280-F15.282, F15.288, F15.29, F16.10, F16.120-F16.122, F16.129, F16.14, F16.150, F16.151, F16.159, F16.180, F16.183, F16.188, F16.19, F16.20, F16.220, F16.221, F16.229, F16.24, 16.250, F16.251, F16.259, F16.280, F16.283, F16.288, F16.29, F18.10, F18.120, F18.121, F18.129, F18.14, F18.150, F18.151, F18.159, F18.17, F18.180, F18.188, F18.19, F18.20, F18.220, F18.221, F18.229, F18.24, F18.250, F18.251, F18.259, F18.27, F18.280, F18.288, F18.29, F19.10, F19.120-F19.122, F19.129, F19.14, F19.150, F19.151, F19.159, F19.16, F19.17, F19.180, F19.181, F19.182, F19.188, F19.19, F19.20, F19.220-F19.222, F19.229-F19.232, F19.239, F19.24, F19.250, F19.251, F19.259, F19.26, F19.27, F19.280-F19.282, F19.288, F19.29

Codes to Identify Outpatient, Intensive Outpatient, Partial Hospitalization, Telehealth, and Medication Treatment Visits (use these visit codes along with the one of the diagnosis codes above to capture initiation and engagement of AOD treatment)

CPT®	HCPCS	UB Revenue
98960-98962, 98966-98969, 99078, 99201-99205, 99211-99215, 99217-99220, 99241-99245, 99281-99285, 99341-99345, 99347-99350, 99384-99387, 99394-99397, 99401-99404, 99408, 99409, 99411, 99412, 99441-99444, 99483, 99510	G0155, G0176, G0177, G0396, G0397, G0409-G0411, G0443, G0463, H0001, H0002, H0004, H0005, H0007, H0008-H0014, H0015,H0016, H0020, H0022, H0031, H0033, H0034-H0037, H0039, H0040, H0047, H2000, H2001, H2010-H2020, H2035, H2036, J0571-J0575, J2315, M0064,S0109, S0201, S9480, S9484, S9485, T1006, T1012, T1015	0100, 0101, 0110-0154, 0156-0160, 0164, 0167, 0169-0174, 0179, 0190-0194, 0199-0204, 0206-0214, 0219, 0450-0452, 0456, 0459, 0510, 0513, 0515-0517, 0519-0523, 0526-0529, 0900, 0902-0907, 0911-0917, 0919, 0944, 0945, 0982, 0981, 0983, 1000-1002
ICD-10		
H2ZZZZ		
CPT WITH POS		Telehealth Modifier
90791, 90792, 90832-90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876	02, 03, 05, 07, 09, 11, 12, 13, 14, 15,16,17,18,19, 20, 22, 33, 49, 50, 52, 53, 57, 71, 72	95, GT
99221-99223, 99231-99233, 99238, 99239, 99251-99255	02, 52, 53	

HOW TO IMPROVE HEDIS SCORES

- ❑ Consider using a validated screening tool (i.e. CAGE-AID, NIDA) or add questions to your standard evaluation to identify substance use issues.
- ❑ Document identified substance abuse in the patient chart and submit a claim with the appropriate codes, as described above.
- ❑ Avoid inappropriate use of diagnosis codes that are the result of alcohol or drug dependency (ex. Cirrhosis) as these also qualify patients for the measures.
- ❑ Schedule a follow-up visit within 14 days and at least two additional visits within 30 days, or refer immediately to a behavioral health provider when giving a diagnosis of alcohol or other drug dependence.
- ❑ Involve family members or others who the patient desires for support and invite their help in intervening with the patient diagnosed with AOD dependence.
- ❑ Provide patient educational materials and resources that include information on the treatment process and options, including 12-Step or peer support meetings, encourage a sponsor and other community-based programs.
- ❑ Refer the member to a Molina Care Manager or work collaboratively with the Molina Care Manager if one is already assigned to help increase member's access and motivation for treatment.
- ❑ Continue ongoing discussions with patients about treatment to help increase their willingness to commit to the process as the timeframe for initiating treatment is 14 days.