

Molina Healthcare

Coding Education

Diabetes Mellitus with Ocular Manifestations



Diabetes is the most common cause of non-congenital vision loss in the United States. There are many ocular complications associated with diabetes including: cataracts, retinopathy, and macular edema.

Cataracts are common and have many etiologies. It is important to document and begin treatment for diabetes mellitus upon diagnosing a diabetic related cataract .

The American Diabetic Association recommends all diabetics undergo an annual comprehensive eye exam by an ophthalmologist or optometrist, including a dilated retinal examination.

Documentation Examples:

Initial Diagnosis

Assessment: 52 year old male with diabetic macular edema, uncontrolled. New worsening of vision, A1C 12.8.

➤ **ICD-10 Code: E11.311** Type 2 diabetes mellitus with unspecified diabetic retinopathy with macular edema and
And

➤ **ICD-10 Code: E11.65** Type 2 diabetes mellitus with hyperglycemia

Plan: Recommend significantly tighter glycemic control, STAT evaluation by ophthalmologist

Established Diagnosis

Assessment: 62 year old male with diabetic proliferative retinopathy and cataract. No macular edema, but noncompliant with recommendations, slight decrease in vision.

➤ **ICD-10 Code: E11.3599** Type 2 diabetes mellitus with proliferative diabetic retinopathy without macular edema, unspecified eye

And

➤ **And ICD-10 Code: E11.36** Type 2 diabetes mellitus with diabetic cataract

Plan: Recommend close follow up with diabetic educator and ophthalmologist/optometrist.

Have Questions?

Contact: Ramp@MolinaHealthcare.com