

<b>Subject: Interspinous Decompression Devices for Spinal Stenosis (X Stop, Coflex)</b>		<b>Original Effective Date:</b> 3/16/2015
<b>Policy Number:</b> MCP-222	<b>Revision Date(s):</b> 7/10/2018	
<b>Review Date:</b> 12/16/15, 9/15/16, 6/22/2017, 6/19/2019		
<b>MCPC Approval Date:</b> 7/10/2018, 6/19/2019		

**DISCLAIMER**

*This Molina Clinical Policy (MCP) is intended to facilitate the Utilization Management process. It expresses Molina's determination as to whether certain services or supplies are medically necessary, experimental, investigational, or cosmetic for purposes of determining appropriateness of payment. The conclusion that a particular service or supply is medically necessary does not constitute a representation or warranty that this service or supply is covered (i.e., will be paid for by Molina) for a particular member. The member's benefit plan determines coverage. Each benefit plan defines which services are covered, which are excluded, and which are subject to dollar caps or other limits. Members and their providers will need to consult the member's benefit plan to determine if there are any exclusion(s) or other benefit limitations applicable to this service or supply. If there is a discrepancy between this policy and a member's plan of benefits, the benefits plan will govern. In addition, coverage may be mandated by applicable legal requirements of a State, the Federal government or CMS for Medicare and Medicaid members. CMS's Coverage Database can be found on the CMS website. The coverage directive(s) and criteria from an existing National Coverage Determination (NCD) or Local Coverage Determination (LCD) will supersede the contents of this Molina Clinical Policy (MCP) document and provide the directive for all Medicare members.<sup>1</sup>*

**DESCRIPTION OF PROCEDURE/SERVICE/PHARMACEUTICAL** <sup>2 3 15 35</sup>

Spinal stenosis is a narrowing of the spinal canal that causes pressure on the spinal cord and nerve roots, low back pain, neurogenic claudication (a combination of low back and leg pain, with numbness and motor weakness when standing or walking), and reduced capacity for physical activity. Symptoms can range from mild to severe, and can affect patient mobility and quality of life. Interspinous decompression devices are intended to be used in patients with lumbar spinal stenosis who have moderately impaired physical function and experience relief in flexion from their symptoms of leg/buttock/groin/back pain, and had failed conservative management. The goal is to provide symptomatic relief of pain, maintain spinal motion, and reduce spine hypermobility and degeneration of adjacent segments levels. There are two types of interspinous devices that include static (i.e X-STOP implant) and dynamic (i.e. non-fusion Coflex). Dynamic devices are intended to be used in conjunction with laminectomy to reduce the amount of lumbar spinal extension possible while preserving range of motion in flexion, axial rotation, and lateral bending. Static devices are used to provide indirect decompression by reducing spinal extension to prevent motions that induce back pain.

The X-Stop Interspinous Spacer device (Medtronic Inc.) was approved by the FDA in 2005 for patient's age > 50 years with moderate symptoms that have not responded adequately to ≥ 6 months of non-operative treatment.<sup>3</sup>

The Coflex Interlaminar Stabilization device (Paradigm Spine LLC) is regulated by the FDA as a spinous process spacer/plate prosthesis that received approval via the premarket approval (PMA) process for treatment of 1- or 2-level LSS from L1-L5 in skeletally mature patients with at least moderately impaired function, relief from buttock/groin/leg pain when in flexion, and 6 months of non-operative treatment.<sup>15</sup>

Vertiflex's Superior® interspinous spacer system received FDA premarket approval in May 2015 for the treatment of moderate stenosis the device is indicated to treat skeletally mature patients suffering from neurogenic intermittent claudication due to moderate degenerative LSS with or without grade 1 spondylolisthesis, who have undergone at least 6 months of non-operative treatment.<sup>35</sup>

## RECOMMENDATION

Interspinous decompression devices (e.g., X-STOP, Coflex, Superior, and any other devices) are considered experimental, investigational and/or unproven for any indication, due to insufficient clinical evidence of safety and efficacy in published peer-reviewed medical literature.

## SUMMARY OF MEDICAL EVIDENCE<sup>2-41</sup>

### X-STOP<sup>2-15</sup>

The clinical evidence of interspinous decompression devices (e.g., X-STOP) for the treatment of lumbar spinal stenosis consists of several randomized controlled trials (RCT's), retrospective comparative studies, prospective studies, and retrospective case series. The study sample sizes ranged from 30 to 285 patients, and follow-up times ranged from 6 months to 4 years. The RCTs compared the efficacy and safety of the device with conservative treatment, decompression, spinal fusion, or with another interspinous spacer. The quality of the overall body of evidence is low and most of the existing studies are small or moderate in size. Additional well-designed, long-term clinical trials are needed to further evaluate the efficacy and safety of interspinous decompression devices and to compare these with standard treatment and other alternatives. A summary of the relevant RCT's is provided below.

One of the earliest RCT's conducted by Zucherman et al. (2004) compared the efficacy and safety of the X Stop with conservative treatment. Results of longer follow-up were reported in a second publication (Zucherman et al., 2005). Participants included 200 patients who had experienced back pain for an average of 4.1 years and who had neurogenic intermittent claudication secondary to LSS. The patients were treated with implantation of 1 or 2 X Stop devices (X Stop group; n=100; mean age 69.9 years) or conservative management with 1 or more epidural steroid injections (Control group; n=91; mean age 68.6 years). Some Control group patients received NSAIDs, analgesics, and/or physical therapy. At 2 years, mean ZCQ symptom severity scores had improved 45% for the X Stop group versus 7% for the Control group. At 1- and 2-year follow-up, there were no significant differences between the 2 groups in any of 8 spinal radiographic measurements. During the 2-year follow-up period, 6% of patients in the X Stop group and 30% in the Control group underwent laminectomy for unresolved symptoms.<sup>7-8</sup>

Another RCT conducted by Azzazi et al. (2010) compared the efficacy and safety of the X Stop with fusion and transpedicular screw fixation in 60 patients with LSS. There were 30 patients in the X Stop group (mean age 57 years, range 28 to 78; mean duration of symptoms 5.4 years), and 30 patients in the Spinal Fusion group (mean age 56.3 years, range 27 to 79; mean duration of symptoms 5.2 years). Outcomes included pain and

disability assessed by the VAS (100-point scale) and the ODI. Patients were followed for 24 months. The source of funding or support for this study was not mentioned. At 24 months, leg pain had decreased significantly from a mean of 82.5 millimeters (mm) preoperatively to 25.5 mm in the X Stop group, and from 80.5 to 35.5 mm in the Spinal Fusion group ( $P<0.01$ ). Back pain improved significantly from 52 mm preoperatively to 29.5 mm in the X Stop group, and from 54 to 37.5 mm in the Spinal Fusion group ( $P<0.01$ ). The ODI score improved significantly from a mean of 53 preoperatively to 26.5 in the X Stop group, and a mean of 55 to 34.5 in the Spinal Fusion group ( $P<0.01$  for these outcomes in each group).<sup>4</sup>

Miller and Block (2012) conducted an RCT that compared the efficacy and safety of the X Stop with the Superior Interspinous Spacer (VertiFlex Inc.), an investigational device that is implanted percutaneously. The X Stop group had 86 patients and the Superior group had 80 patients (mean age 67 years in each group). Data from the first 6 months of assessment were available for 30 patients in the X Stop group and 36 patients in the Superior group. By 6 months, ZCQ symptom severity scores improved by 25% and 30%, and physical function scores improved by 27% and 32% in the X Stop and Superior groups, respectively ( $P<0.001$  for all analyses). The proportion of patients who had ZCQ clinical success was 53% and 75% for symptom severity, 63% and 64% for physical function, and 93% and 78% for patient satisfaction in the X Stop and Superior groups, respectively. By 6 months, axial pain improved by 64% and 70%, and extremity pain improved by 81% and 93% in the X Stop and Superior groups, respectively ( $P<0.001$ ). Clinical success for axial pain was achieved by 50% and 60%, and clinical success for extremity pain by 60% and 74% in the X Stop and Superior groups, respectively. By 6 months, back function improved by a median of 38% and 48%, and back function clinical success was achieved by 37% and 47% in the X Stop and Superior groups, respectively ( $P<0.001$  for all analyses). The authors reported on significant improvements in pain and disability within each treatment group, but did not systematically compare outcomes for patients treated with the X Stop and the Superior devices.<sup>5</sup>

Another RCT conducted by Strömquist et al. (2013) compared the efficacy and safety of the X Stop with standard decompressive surgery in 100 patients with LSS. There were 50 patients in the X Stop group (mean age 67 years, range 49 to 89) and 50 patients in the Decompression group (mean age 71 years, range 57 to 84); all patients had symptoms of neurogenic claudication for at least 6 months. Outcomes included pain and disability assessed by the VAS, ZCQ, ODI, SF-36, and European Quality of Life 5 dimension questionnaire (EQ-5D; EuroQol Group). Patients were evaluated at 6 months, 12 months, and 2 years. Symptom severity and physical function according to ZCQ were significantly improved by 6 months in both groups. The SF-36 physical component score improved from 25 and 28 points preoperatively to 40 and 38 points at 2 years postoperatively, for the X Stop and Decompression groups, respectively. As measured by VAS, mean back pain improved from 58 and 60 points preoperatively to 34 and 23 points at 2 years postoperatively, for the X Stop and Decompression groups, respectively. Mean left leg pain improved from 57 and 58 points preoperatively to 25 and 19 points at 2 years postoperatively and mean right leg pain improved from 60 and 53 points preoperatively to 21 and 21 points postoperatively, respectively. The changes in pain from baseline were significant in both groups ( $P<0.001$ ); however, there were no significant differences between groups.<sup>6</sup>

### Coflex<sup>16-33</sup>

The clinical evidence of interspinous decompression devices (e.g., Coflex) for the treatment of lumbar spinal stenosis consists of RCT's, prospective nonrandomized comparative study, retrospective comparative studies, and retrospective case series that evaluated the efficacy and safety of the Coflex interlaminar stabilization

device for treating symptomatic lumbar spinal stenosis. The study sample sizes ranged from 20 to 322 patients, and mean follow-up ranged from 11.8 months to 6.3 years. Outcomes included pain levels and function assessed by a VAS or other scale for pain; narcotics use, ROM, and/or neurological examination. The quality of the overall body of evidence is low and most of the existing studies are small or moderate in size. Additional well-designed, long-term clinical trials are needed to further evaluate the efficacy and safety of interspinous decompression devices and to compare these with standard treatment and other alternatives. A summary of the relevant RCT's is provided below.

A moderate size RCT (Davis et al. (2013a) compared the efficacy and safety of spinal decompression plus coflex with decompression plus fusion in 322 patients with LSS, and in a subset of 150 patients with grade I spondylolisthesis. Both treatments led to significant improvement at 24 months in mean scores on the VAS for back pain and leg pain, ODI, SF-12 physical component, and ZCQ symptom severity and physical function, compared with baseline values. At 24 months, mean scores for the SF-12 physical component and ZCQ symptom severity, physical function, and patient satisfaction were significantly better for the coflex than for fusion; however, mean VAS and ODI scores were similar for the 2 approaches in the entire cohort. In the entire cohort and the in the subset with spondylolisthesis, the mean SF-12 mental component score did not change appreciably and was similar between the Coflex and Fusion groups at all evaluation times. At 24 months, radiographic results revealed changes in ROM in patients who had fusion (rotation and translation decreased at the treated lumbar level(s) and increased at the level above and the level below the treated level(s)). In contrast, ROM was fairly well preserved (rotation and translation changed by  $< 1.0^\circ$  or  $< 1.0$  mm, respectively, at treated and adjacent levels) in the Coflex group.<sup>19</sup>

In a secondary analysis of this RCT, Davis et al. (2013b) reported on the outcomes of a subset of 150 patients with Meyerding grade I spondylolisthesis ( $\leq 25\%$  sagittal plane translation on flexion-extension radiographs) who were included in the randomized FDA IDE trial in the Coflex group (n=99; mean age 63.1 years, range not reported; 41% men; 2-level procedures required in 64.2%; mean ODI 59.4; mean VAS for back pain 80.3; mean VAS for worse leg pain 77.9) or the Fusion group (n=51; mean age 65.0 years, range not reported; 19% men; 2-level procedures required in 63.6%; mean ODI 60.0; mean VAS for back pain 78.6; mean VAS for worse leg pain 79.1). Follow-up findings were reported only for the 24-month evaluation, at which time data were available in 94.9% of the Coflex group and 94.1% of the Fusion group. Both groups demonstrated significant improvement in mean scores for ODI ( $-38.3$  and  $-37.1$  points, respectively), VAS for back pain ( $-54.9$  and  $-58.0$  mm), VAS for worse leg pain ( $-58.9$  and  $-56.2$  points), SF-12 physical component (16.4 and 14.8 points), ZCQ symptom severity ( $-1.64$  and  $-1.40$  points), and ZCQ physical function ( $-1.24$  and  $-1.10$  points). The rate of composite clinical success was similar in the Coflex and Fusion groups (62.8% and 62.5%, respectively).<sup>20</sup>

Bae et al. (2016) performed a three year follow-up analysis of the Davis (2013a) RCT. At 36 months, 91% (195/215) of the coflex group and 88% (94/107) of the fusion group were included in the analysis. The initial efficacy endpoints (composite scores) were modified for use at 36 months. At 36 months, 62.2% of the individuals in the coflex group compared to 48.9% of the individuals in the 94 group reported composite clinical success scores (difference = 13.3%, 95% confidence interval [CI]; 1.1%-25.5%,  $p=0.03$ ). There are several limitations in this study including the limited follow-up period and the heterogeneous mix of individuals including those without listhesis for which fusion/stabilization is an unproven procedure.<sup>24</sup>

Published reports of the Superion Interspinous Spacer includes a randomized controlled trial, and prospective case series. Although the randomized device trials report short-term improvements in symptoms and functional status when compared to non-operative therapy, a number of questions remain. Overall, high-quality comparative data are limited and there is a lack of evidence to support that interspinous spacer devices are as safe and effective as the gold standard of decompression. In addition, there appears to be some concerns that the devices are not as effective as surgical decompression and lead to higher rates of reoperation.

Patel et al. (2015b) published a report on 3-year durability of results of the pivotal trial. At 36 months, the overall treatment success (primary composite endpoint) remained stable in the Superion group (52.5% of 120 participants available for follow-up at 36 months versus 52.7% at 24 months). In the X-Stop group, the composite endpoint of overall treatment success was 38.0% of 129 participants available for follow-up at 36 months, reduced from 50.2% at 24 months. The difference between groups was statistically significant (P=0.023). A total of 26 (14%) participants in the Superion group required surgical decompression within 3 years. A majority of patients in the Superion group experienced significant improvements in individual outcome measures, including back pain as measured by a  $\geq 20$  mm decrease in visual analog scale (VAS) (76.8%), VAS leg pain (84.1%), ZCQ physical function (80.5%), ZCQ symptom severity (82.9%), Oswestry Disability Index ( $\geq 15$ -point decrease) (69.5%), and ZCQ patient satisfaction (91.5%) at 36 months. Between-group differences in most individual outcome measures were not statistically significant, with the exception of VAS leg pain. A total of 69.7% of patients in the X-Stop group had durable improvement in leg pain at 36 months, compared with 84.1% of the Superion group (P=0.037).<sup>35</sup>

Nunley et al (2017) reported five-year clinical outcomes of a randomized controlled U.S. FDA noninferiority trial in individuals with moderate lumbar spinal stenosis. While the original trial compared the Superion to the X STOP device, the analysis was restricted to the Superion trial arm. A total of 73% (88/121) of the living individuals who received the spacer device participated in the 5 year clinical outcomes assessment. Outcomes were assessed using the Zurich Claudication Questionnaire (ZCQ), leg and back pain severity by visual analog scale (VAS), and the Oswestry Disability Index (ODI). The authors reported success rates in all areas of assessment, 84% reported clinical success in at least two of the three ZCQ domains, 80% leg pain VAS scores, 65% back pain VAS scores and 65% for ODI scores. There remains a lack of studies which compare interspinous spacers to standard treatments, such as decompression surgery.<sup>38</sup>

**CODING INFORMATION** THE CODES LISTED IN THIS POLICY ARE FOR REFERENCE PURPOSES ONLY. LISTING OF A SERVICE OR DEVICE CODE IN THIS POLICY DOES NOT IMPLY THAT THE SERVICE DESCRIBED BY THIS CODE IS COVERED OR NON-COVERED. COVERAGE IS DETERMINED BY THE BENEFIT DOCUMENT. THIS LIST OF CODES MAY NOT BE ALL INCLUSIVE.

CPT	Description
22867	Insertion of interlaminar/interspinous process stabilization/distraction device, without fusion, including image guidance when performed, with open decompression, lumbar; single level
22868	Insertion of interlaminar/interspinous process stabilization/distraction device, without fusion, including image guidance when performed, with open decompression, lumbar; second level (List separately in addition to code for primary procedure)

22869	Insertion of interlaminar/interspinous process stabilization/distraction device, without open decompression or fusion, including image guidance when performed, lumbar; single level
22870	Insertion of interlaminar/interspinous process stabilization/distraction device, without open decompression or fusion, including image guidance when performed, lumbar; second level (List separately in addition to code for primary procedure)

HCPCS	Description
C1821	Interspinous process distraction device (implantable)

ICD-10	Description: [For dates of service on or after 10/01/2015]
M48.06-M48.07	Spinal stenosis

## REFERENCES

### *X-STOP*

- Centers for Medicare & Medicaid Services (CMS). Medicare Coverage Database. Advanced Search: National Coverage Documents [search:]. Available at: <http://www.cms.gov/medicare-coverage-database/>
- Center for Devices and Radiological Health (CDRH). Premarket Approval (PMA) Database. Food and Drug Administration [website]. Search P040001 in the P Number field. Available at: <http://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfPMA/pma.cfm>
- Hayes Health Technology Brief. X Stop Interspinous Process Decompression System (Medtronic Spine LLC) for Lumbar Spinal Stenosis. Winifred Hayes Inc. Lansdale, Pa. Feb, 2014. **Archived Jan 2016.**
- Azzazi A, Elhawary Y. Dynamic stabilization using X-stop versus transpedicular screw fixation in the treatment of lumbar canal stenosis; comparative study of the clinical outcome. *Neurosurg Q.* 2010;20(3):165-169.
- Miller LE, Block JE. Interspinous spacer implant in patients with lumbar spinal stenosis: preliminary results of a multicenter, randomized, controlled trial. *Pain Res Treat.* 2012;2012:823509. Available at: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3289933>. Accessed February 24, 2014.
- Strömqvist BH, Berg S, Gerdhem P, et al. X-Stop versus decompressive surgery for lumbar neurogenic intermittent claudication: randomized controlled trial with 2-year follow-up. *Spine (Phila Pa 1976).* 2013;38(17):1436-1442.
- Zucherman JF, Hsu KY, Hartjen CA, et al. A multicenter, prospective, randomized trial evaluating the X STOP interspinous process decompression system for the treatment of neurogenic intermittent claudication: two-year follow-up results. *Spine (Phila PA 1976).* 2005;30(12):1351-1358.
- Zucherman JF, Hsu KY, Hartjen CA, et al. A prospective randomized multi-center study for the treatment of lumbar spinal stenosis with the X STOP interspinous implant: 1-year results. *Eur Spine J.* 2004;13(1):22-31. Available at: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3468027>. Accessed February 24, 2014.

9. Puzzilli F, Gazzeri R et al. Interspinous spacer decompression (X-STOP) for lumbar spinal stenosis and degenerative disk disease: a multicenter study with a minimum 3-year follow-up. *Clin Neurol Neurosurg*. 2014 Sep;124:166-74. doi: 10.1016/j.clineuro.2014.07.004. Epub 2014 Jul 14.
10. Wu AM, Zhou Y et al. Interspinous spacer versus traditional decompressive surgery for lumbar spinal stenosis: a systematic review and meta-analysis. *PLoS One*. 2014 May 8;9(5):e97142. doi: 10.1371/journal.pone.0097142. eCollection 2014.
11. Patel VV, Whang PG et al. Two-year clinical outcomes of a multicenter randomized controlled trial comparing two interspinous spacers for treatment of moderate lumbar spinal stenosis. *BMC Musculoskelet Disord*. 2014 Jul 5;15:221. doi: 10.1186/1471-2474-15-221.
12. Hartjen CA1, Resnick DK et al. Two-Year Evaluation of the X-STOP Interspinous Spacer in Different Primary Patient Populations With Neurogenic Intermittent Claudication due to Lumbar Spinal Stenosis. *J Spinal Disord Tech*. 2013 Nov 8. [Epub ahead of print]
13. Lonne G, et al. Minimally invasive decompression versus x-stop in lumbar spinal stenosis: a randomized controlled multicenter study. *Spine* 2015 Jan 15;40(2):77-85.
14. Lonne G, et al. Comparing cost-effectiveness of x-stop with minimally invasive decompression in lumbar spinal stenosis: A randomized controlled trial. *Spine* 2015 Apr 15;40(8):514-20.
15. Gazzeri R, Galarza M et al. Failure rates and complications of interspinous process decompression devices: a European multicenter study. *Neurosurg Focus*. 2015 Oct;39(4):E14.

### **Coflex**

16. Hayes Health Technology Brief. Coflex Interlaminar Stabilization Device (Paradigm Spine LLC) for Treatment of Lumbar Spinal Stenosis. Winifred Hayes Inc. Lansdale, Pa. **Sept, 2018**.
17. Center for Devices and Radiological Health (CDRH). Premarket Approval (PMA) Database. Food and Drug Administration [website]. Available at: [http://www.accessdata.fda.gov/cdrh\\_docs/pdf11/P110008a.pdf](http://www.accessdata.fda.gov/cdrh_docs/pdf11/P110008a.pdf)
18. Cabraja M, Abbushi A, Woiciechowsky C, Kroppenstedt S. The short- and mid-term effect of dynamic interspinous distraction in the treatment of recurrent lumbar facet joint pain. *Eur Spine J*. 2009;18(11):1686-1694. Available at: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2899396>.
19. Davis RJ, Errico TJ, Bae H, Auerbach JD. Decompression and Coflex interlaminar stabilization compared with decompression and instrumented spinal fusion for spinal stenosis and low-grade degenerative spondylolisthesis. two-year results from the prospective, randomized, multicenter, Food and Drug Administration Investigational Device Exemption trial. *Spine (Phila Pa 1976)*. 2013a;38(18):1529-1539.
20. Davis R, Auerbach JD, Bae H, Errico TJ. Can low-grade spondylolisthesis be effectively treated by either coflex interlaminar stabilization or laminectomy and posterior spinal fusion? Two-year clinical and radiographic results from the randomized, prospective, multicenter US investigational device exemption trial: clinical article. *J Neurosurg Spine*. 2013b;19(2):174-184. Available at: <http://thejns.org/doi/full/10.3171/2013.4.SPINE12636>.
21. Park SC, Yoon SH, Hong YP, Kim KJ, Chung SK, Kim HJ. Minimum 2-year follow-up result of degenerative spinal stenosis treated with interspinous U (Coflex™). *J Korean Neurosurg Soc*. 2009;46(4):292-299. Available at: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2773383>.

22. Richter A, Halm HF, Hauck M, Quante M. 2-year follow-up after decompressive surgery with and without implantation of an interspinous device for lumbar spinal stenosis: a prospective controlled study. *J Spinal Disord Tech.* 2014 Aug;27(6):336-41. doi: 10.1097/BSD.0b013e31825f7203.
23. Kumar N, Shah SM et al. Role of coflex as an adjunct to decompression for symptomatic lumbar spinal stenosis. *Asian Spine J.* 2014 Apr;8(2):161-9. doi: 10.4184/asj.2014.8.2.161. Epub 2014 Apr 8.
24. Bae HW, Davis RJ, Lauryssen C, Leary S, Maislin G, Musacchio M Jr. Three-year follow-up of the prospective, randomized, controlled trial of coflex interlaminar stabilization vs instrumented fusion in patients with lumbar stenosis. *Neurosurgery.* 2016. Epub ahead of print. April 4, 2016. Available at: <http://journals.lww.com/neurosurgery/pages/articleviewer.aspx?year=9000&issue=00000&article=97361&type=abstract>
25. Bae HW, Lauryssen C, Maislin G, Leary S, Musacchio MJ Jr. Therapeutic sustainability and durability of coflex interlaminar stabilization after decompression for lumbar spinal stenosis: a four year assessment. *Int J Spine Surg.* 2015;9:15.
26. Musacchio MJ, Lauryssen C, Davis RJ, et al. Evaluation of decompression and interlaminar stabilization compared with decompression and fusion for the treatment of lumbar spinal stenosis: 5-year follow-up of a prospective, randomized, controlled trial. *Int J Spine Surg.* 2016;10:6.
27. Li AM, et al. Decompression and coflex interlaminar stabilization compared with conventional surgical procedures for lumbar spinal stenosis: A systematic review and meta-analysis. *Int J Surg* 2017 April;40:60-67.
28. Xu C, et al. Application of the Coflex interlaminar stabilization in patients with L5/S1 degenerative diseases: minimum 4-year follow-up. *Am J Ther* 2016 Nov/Dec;23(6):e1813-e1818.
29. Kamal TT, et al. Reported clinical outcomes of coflex dynamic stabilization device vs instrumented decompression and fusion in degenerative lumbar stenosis. *MOJ Orthoped Rheumat* 2016;5(6):00200.
30. Lee N, et al. Paradoxical radiographic changes of Coflex Interspinous Device with minimum 2-year follow-up in lumbar spinal stenosis. *World Neurosurg* 2016 Jan;85:177-184.
31. Abjornson C, Yoon BJV, Callanan T, Shein D, Grinberg S, Cammisa FP. Spinal stenosis in the absence of spondylolisthesis: can interlaminar stabilization at single and multi-levels provide sustainable relief? *Int J Spine Surg.* 2018;12(1):64-69.
32. Schmidt S, Franke J, Rauschmann M, Adelt D, Bonsanto MM, Sola S. Prospective, randomized, multicenter study with 2-year follow-up to compare the performance of decompression with and without interlaminar stabilization. *J Neurosurg Spine.* 2018;28(4):406-415.
33. Zhang JX, Jing XW, Cui P, He X, Hao DJ, Li SJ. Effectiveness of dynamic fixation coflex treatment for degenerative lumbar spinal stenosis. *Exp Ther Med.* 2018;15(1):667-672.

## Superion

34. Patel VV, et al. Superion interspinous process spacer for intermittent neurogenic claudication secondary to moderate lumbar spinal stenosis: two-year results from a randomized controlled FDA-IDE pivotal trial. *Spine* 2015a Mar 1;40(5):275-82.
35. Patel VV, Nunley PD, Whang PG et al. Superion(®) InterSpinous Spacer for treatment of moderate degenerative lumbar spinal stenosis: durable three-year results of a randomized controlled trial. *J Pain Res.* 2015b Oct 3;8:657-62. doi: 10.2147/JPR.S92633. eCollection 2015.

36. Patel VV, et al. Superior® Interspinous Spacer for treatment of moderate degenerative lumbar spinal stenosis: durable three-year results of a randomized controlled trial. *J Pain Res* 2015 Oct 3;8:657-62.
37. Nunley PD, Patel VV, Orndorff DG, et al. Superior Interspinous Spacer Treatment of Moderate Spinal Stenosis: 4-Year Results. *World Neurosurg*. 2017 Aug;104:279-283. doi: 10.1016/j.wneu.2017.04.163. Epub 2017 May 4.
38. Nunley PD, Patel VV, Orndorff DG et al. Five-year durability of stand-alone interspinous process decompression for lumbar spinal stenosis. *Clin Interv Aging*. 2017 Sep 6;12:1409-1417. doi: 10.2147/CIA.S143503. eCollection 2017.
39. Hayes. Winifred Hayes, Inc. Lansdale, PA.
  - Prognosis Overview. Superior Interspinous Spacer System. Feb, 2015. Archived Aug, 2016.
  - CRR. Superior Interspinous Spacer System (Vertiflex Inc.) for Spinal Stenosis. May, 2018.
40. Zaina F, Tomkins-Lane C, Carragee E, Negrini S. Surgical versus non-surgical treatment for lumbar spinal stenosis. *Cochrane Database Syst Rev*. 2016 Jan 29;(1):CD010264
41. Zhao H, Duan LJ, Gao YS, Yang YD et al. Comparison of two FDA-approved interspinous spacers for treatment of lumbar spinal stenosis: Superior versus X-STOP-a meta-analysis from five randomized controlled trial studies. *J Orthop Surg Res*. 2018 Mar 2;13(1):42.

## Other Resources

42. Official Disability Guidelines: Interspinous decompression device. 2015.
43. Guyer R, et al. ISASS Recommendations/coverage criteria for decompression with interlaminar stabilization-coverage indications, limitations, and/or medical necessity. *Int J Spine Surg* 2016 Dec 5;10:41. Accessed at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5374995/>
44. UpToDate: [Website] Waltham, MA: Walters Kluwer Health; 2019
  - Levin K. Lumbar spinal stenosis: Treatment and prognosis.
45. Advanced Medical Review: Policy reviewed by a practicing physician board certified in Orthopaedic Surgery. 3/27/18.

## Professional Society Guidelines

46. American Association of Neurological Surgeons (AANS) Lumbar Spinal Stenosis. 2018. Available at: <http://www.aans.org/en/Patients/Neurosurgical-Conditions-and-Treatments/Lumbar-Spinal-Stenosis>.
47. Ghiselli G, Kreiner S. North American Spine Society (NASS) coverage policy recommendations. Lumbar Interspinous Device without Fusion and with Decompression. NASS Coverage Committee. May, 2018.
48. Guyer R, Musacchio M, Lorio MP. International Society for the Advancement of Spine Surgery (ISASS) recommendations/coverage criteria for decompression with interlaminar stabilization - coverage indications, limitations, and/or medical necessity. *Int J Spine Surg*. 2016;10:41.

## Review/Revision History:

3/16/15: Policy created

12/16/15, 9/15/16, 6/22/17: Policy reviewed, no changes.

7/10/18: This policy was reviewed and the clinical criteria has not changed, these devices remains controversial and experimental for the treatment of spinal stenosis. The following sections were updated: new device called

Vertiflex's Superior® interspinous spacer system was added into the description section, summary of medical evidence, professional society guidelines, CPT coding tables and references were updated.

6/19/19: This policy was reviewed and the clinical criteria has not changed, these devices remains controversial and experimental for the treatment of spinal stenosis. The following sections were updated: professional society guidelines and references.