

Molina Clinical Policy

Transarterial Chemoembolization (TACE) and Transarterial Embolization (TAE) for Liver Tumors: Policy No. 120

Last Approval: 4/5/2021

Next Review Due By: April 2022



DISCLAIMER

This Molina Clinical Policy (MCP) is intended to facilitate the Utilization Management process. Policies are not a supplementation or recommendation for treatment; Providers are solely responsible for the diagnosis, treatment and clinical recommendations for the Member. It expresses Molina's determination as to whether certain services or supplies are medically necessary, experimental, investigational, or cosmetic for purposes of determining appropriateness of payment. The conclusion that a particular service or supply is medically necessary does not constitute a representation or warranty that this service or supply is covered (e.g., will be paid for by Molina) for a particular Member. The Member's benefit plan determines coverage – each benefit plan defines which services are covered, which are excluded, and which are subject to dollar caps or other limits. Members and their Providers will need to consult the Member's benefit plan to determine if there are any exclusion(s) or other benefit limitations applicable to this service or supply. If there is a discrepancy between this policy and a Member's plan of benefits, the benefits plan will govern. In addition, coverage may be mandated by applicable legal requirements of a State, the Federal government or CMS for Medicare and Medicaid Members. CMS's Coverage Database can be found on the CMS website. The coverage directive(s) and criteria from an existing National Coverage Determination (NCD) or Local Coverage Determination (LCD) will supersede the contents of this MCP and provide the directive for all Medicare members.¹ References included were accurate at the time of policy approval and publication.

OVERVIEW ^{34,35,42,43}

Transarterial chemoembolization (TACE) or Transarterial embolization (TAE) are catheter-based embolization procedures. TACE is combined with a prior injection of a chemotherapeutic agent into the hepatic artery, and TAE is not combined with a chemotherapeutic agent (bland embolization). Both embolization procedures lead to ischemia of the tumor by blockage of the nutrient supply. These procedures have been investigated to treat resectable, unresectable, and recurrent HCC, as a bridge to liver transplantation, and to treat liver metastases, most commonly from colorectal cancer. These procedures are a two-step process that involves placing a catheter in the artery that is supplying blood to a tumor to precisely infuse first, a chemotherapeutic agent and then, an embolic agent. This process effectively delivers a drug and blocks blood flow to the tumor. TAE is proposed as a palliative treatment for primary hepatic or metastatic tumors including neuroendocrine tumors with hepatic metastases. TAE is conducted during selective hepatic arterial catheterization through the arteries supplying a tumor and involves the infusion of lipiodol (without a chemotherapeutic agent) followed by embolization using any of the embolic agents (e.g., gelatin sponge cubes) as during TACE procedures. TACE and TAE require hospitalization and multiple treatments may be required to treat all lesions as well as recurrences, however, the benefit of repetition needs to be balanced against the progressive liver damage associated with the treatment. The most common adverse effect of TACE and TAE are post-embolization syndrome which consists of varying degrees of right upper quadrant pain, nausea, a moderate degree of ileus, fatigue, fever, and transient elevation of AST, ALT and bilirubin values. Symptoms are usually self-limited, lasting three to four days; full recovery is typical within 7 to 10 days.

Drug-eluting beads transarterial chemoembolization (DEB-TACE) is an adaptation of TACE and is a single-step process where beads of uniform size are loaded with a drug and delivered to the feeder artery in a single step. This procedure is easier to perform than two-step TACE. Since the drug is contained within the beads, its elution is more localized and longer lasting than in traditional TACE. Embolic beads, or microspheres, are either synthetic polymers such as ethylene vinyl acetate, or natural materials such as albumin, chitosan, gelatin, or alginate. Doxorubicin is the most commonly used chemotherapeutic agent for DEB-TACE, however mitomycin C, cisplatin, methotrexate, and paclitaxel are also used. DEB-TACE is prescribed by an oncologist and performed in an angiography suite by an interventional radiologist. The procedure takes approximately 2 hours and patients are hospitalized for 1 to 2 days. Bilateral disease is treated in separate procedures at an interval of approximately 2 weeks. Repeat DEB-TACE procedures may be conducted, up to 4 times over a 6-month period.²⁵ Embosphere Microspheres are spherical, tightly calibrated, biocompatible, nonresorbable, hydrogel microspheres coated with an inorganic perfluorinated polymer (Polyzene-F) and are used during transarterial embolization (TAE) to decrease the blood supply of unresectable, intermediate- and advanced-stage hepatocellular carcinoma (HCC).

Hepatocellular carcinoma (HCC) is a primary tumor of the liver that usually develops in the setting of chronic liver disease, particularly in patients with chronic hepatitis B and C. TACE is an appropriate option for patients with a large unresectable or multifocal HCC without main or lobar branch portal vein thrombus that is not amenable to local ablation. TACE is also commonly used as a bridging maneuver in patients awaiting liver transplantation. For most patients who are candidates for resection, preoperative TACE is not indicated.

Molina Clinical Policy
Transarterial Chemoembolization (TACE) and Transarterial Embolization (TAE) for Liver Tumors: Policy No. 120

Last Approval: 4/5/2021
Next Review Due By: April 2022



Neuroendocrine tumors (NETs) are a heterogeneous group of neoplasms that are thought to arise from neuroendocrine cells and their precursors located throughout the body. TACE is applied as a palliative technique in patients with a hepatic-predominant metastatic NET who are not candidates for surgical resection.

Uveal melanoma is a rare malignancy that arises from melanocytes within the uveal tract of the eye, which includes the iris, ciliary body, and choroid. Uveal melanoma comprises approximately 95 percent of melanomas arising from the eye, with the remainder arising from the conjunctiva. Among patients with hepatic metastases, TACE directed specifically toward the liver metastases has been associated with responses that may have clinical utility.

Colorectal Cancer gold standard in management of colon cancer metastatic to the liver remains resection. However, most patients are not candidates for surgery due to either disease bulk or the presence of extrahepatic metastases. Arterial therapies such as TACE and SIRT, either as monotherapy or in combination with other therapeutic regimens, have shown survival benefit.

COVERAGE POLICY 2-15,24-33,36-41,44,45

Transarterial chemoembolization (TACE) or Transarterial embolization (TAE) **is considered medically necessary** for any of the following conditions:

1. Treatment of primary hepatocellular liver carcinoma (HCC) or as a bridge to liver transplant when **ALL** of the following criteria are met:
 - Preserved liver function defined as Childs-Turcotte-Pugh Class A or B; **and**
 - Localized unresectable or multifocal tumor with **ALL** of the following:
 - Three or fewer encapsulated nodules; **AND**
 - Each nodule is less than or equal to 5 centimeters in diameter.

AND

 - No evidence of extra-hepatic metastases; **AND**
 - No evidence of severe renal function impairment; **AND**
 - No evidence of portal vein occlusion.

OR

2. Treatment of primary HCC³⁶⁻⁴¹ in individuals who may become eligible for liver transplantation when the following criteria are met:
 - a. One lesion greater than 5 cm and less than or equal to 8 cm; **OR**
 - b. Two or three lesions each greater than 3 cm and less than or equal to 5 cm; **AND**
 - c. A total diameter of all lesions less than or equal to 8 cm; **OR**
 - d. Four to five lesions each less than 3 cm, and a total diameter of all lesions less than or equal to 8 cm.

OR

3. Treatment of liver metastasis in symptomatic patients with metastatic neuroendocrine tumors^{36 44 45} whose symptoms persist despite systemic treatment and who are not candidates for surgical resection; **OR**
4. Treatment of liver metastasis in patients with liver-dominant metastatic uveal melanoma;³⁶⁻⁴¹ **OR**
5. Treatment of liver metastasis in select patients with colorectal cancer whose symptoms persist despite systemic treatment and who are not candidates for surgical resection.^{36 44}

Molina Clinical Policy

Transarterial Chemoembolization (TACE) and Transarterial Embolization (TAE) for Liver Tumors: Policy No. 120

Last Approval: 4/5/2021

Next Review Due By: April 2022



* The Child-Turcote-Pugh (CPT) score determines short-term prognosis among groups of patients awaiting liver transplantation and has been widely adopted for risk-stratifying patients before transplantation.

Child-Turcote-Pugh Score of Severity of Liver Disease ³⁴			
Points	1	2	3
Encephalopathy	None	1 – 2	3 – 4
Ascites	Absent	Slight	Moderate
Bilirubin (mg/dL)	< 2	2 – 3	> 3
For PBC/PSC, Bilirubin	< 4	4 – 10	> 10
Albumin (g/dL)	> 3.5	2.8 – 3.5	< 2.8
INR*	< 1.7	1.7 – 2.3	> 2.3
PT (seconds prolonged)	< 4	4 – 6	> 6

The individual scores are summed and then grouped as a classification: < 7 = A, 7-9 = B, > 9 = C (forecasts a survival of less than 12 months). *INR = International Normalized Ratio; PT = prothrombin time.

Continuation of Therapy⁴²

The TACE procedure may be repeated if there is clear evidence of progressive tumor growth in the treated areas.

- Multiple courses of TACE, especially if spaced too closely together, can increase deaths from liver failure despite successful tumor shrinkage, and these excess deaths from deterioration of liver function may counterbalance any prolongation of survival that results from enhanced tumor control.
- TACE may cause hepatic artery damage, the likelihood of which is higher in patients with impaired liver function.
- Hepatic artery interruption by repeated TACE or arterial dissection also leads to the development of extrahepatic collateralization, which may create an alternative blood supply to the tumor and contribute to treatment failure.

Limitations and Exclusions^{36,37,42}

1. TACE utilizing chemotherapy-loaded microspheres (e.g., drug-loaded microspheres, drug-eluting beads, and doxorubicin drug-eluting bead transarterial chemoembolization [DEB-TACE] and Embozene Microspheres) **are considered experimental, investigational and unproven** for all liver-related conditions.
2. TACE is contraindicated for **ANY** of the following conditions:
 - Absent or severely reduced portal vein flow (eg, tumoral or nontumoral portal vein occlusion, or hepatofugal blood flow); **AND**
 - Decompensated cirrhosis (Child-Turcotte-Pugh C, or Child-Turcotte-Pugh B score >8 including jaundice, clinical hepatic encephalopathy, refractory ascites, and/or hepatorenal syndrome).
3. Relative contraindications include **ANY** of the following:
 - Serum bilirubin >2 mg/dL; **OR**
 - Lactate dehydrogenase >425 units/L; **OR**
 - Aspartate aminotransferase >100 units/L; **OR**
 - Tumor burden involving >50 percent of the liver; **OR**
 - Severe comorbidities; **OR**
 - Untreated esophageal varices at high risk of bleeding; **OR**
 - Prior transjugular intrahepatic portosystemic shunting (TIPS).

DOCUMENTATION REQUIREMENTS. Molina Healthcare reserves the right to require that additional documentation be made available as part of its coverage determination; quality improvement; and fraud; waste and abuse prevention processes. Documentation required may include, but is not limited to, patient records, test results and credentials of the provider ordering or performing a drug or service. Molina Healthcare may deny reimbursement or take additional appropriate action if the documentation provided does not support the initial determination that the drugs or services were medically necessary, not investigational or experimental, and otherwise within the scope of benefits afforded to the member, and/or the documentation demonstrates a pattern of billing or other practice that is inappropriate or excessive.

SUMMARY OF MEDICAL EVIDENCE

The available evidence on the efficacy and safety of DEB-TACE with doxorubicin for unresectable HCC is conflicting and consists of RCTs (n=30-201), that compared the efficacy and safety of treatment with DEB-TACE

Molina Clinical Policy

Transarterial Chemoembolization (TACE) and Transarterial Embolization (TAE) for Liver Tumors: Policy No. 120

Last Approval: 4/5/2021

Next Review Due By: April 2022



with conventional TACE or compared it with bland embolization with inert beads. Additional published evidence consists of controlled retrospective, uncontrolled, and prospective trials (n=62-71). The overall quality of the evidence is low because of study size, limitations of the uncontrolled studies, and the lack of RCTs. The available evidence suggests that DEB-TACE with doxorubicin is a safe and efficacious treatment for unresectable HCC, however there is insufficient evidence to draw conclusions regarding superiority of DEB-TACE over conventional TACE. Additional trials are necessary to compare the efficacy and safety of DEB-TACE with conventional TACE, assess quality of life and assess a consistent imaging method to monitor the tumor response to DEB-TACE.¹⁶⁻²³

The available evidence on the efficacy and safety of TACE or TAE to treat unresectable HCC is from limited RCT's, (n=40-400) systematic reviews, retrospective reviews, and prospective studies (n=120-1293). Overall survival rates for chemoembolization as palliative treatment of unresectable disease ranged from 42% to 93% at 1 year, 22% to 87% at 2 years, and 3% to 43% at 3 years. Although the data is limited it does demonstrate that TACE provides a survival benefit for the treatment of unresectable primary hepatocellular carcinoma (HCC) when compared with supportive care only or when added as an additional therapy versus supportive care or systemic chemotherapy alone in the management of unresectable HCC.²⁻¹⁵ Consensus opinion from professional organizations suggests that TACE may be a treatment option for unresectable HCC.³⁶⁻⁴¹

The available evidence on the efficacy and safety of TACE or TAE as a bridge to liver transplant consists of a number of uncontrolled studies that report that TACE is associated with low rates of dropout from the transplant list, and is likely to reduce dropouts from the list. As a result, TACE has become an accepted component of care for patients with HCC on the waiting list for liver transplant. The available evidence on the efficacy and safety of TACE for patients with unresectable neuroendocrine tumors consists of uncontrolled trials and case studies that have reported that TACE reduces symptoms and tumor burden, improves hormone profile and a prolonged hepatic progression-free survival. Reports state that TACE generally achieves average symptomatic, biological and radiological responses of 75%, 56% and 50%, with progression-free survival of 12-18 months, with acceptable tolerance. In addition, for its relatively rare condition, there are limited alternative treatments for these tumors. The available evidence on the efficacy and safety of TACE for hepatic metastases from uveal (ocular) melanoma includes retrospective studies, case series of treated patients that have reported that tumor response and survival are improved compared to historical controls. The treatment with TACE conferred a survival advantage (median 16.5 vs. 12.2 months for other first line treatments). There are limited treatment options and this condition is rare, making the performance of high-quality RCTs difficult or impossible. These reports conclude that TACE improves outcomes for patients with hepatic metastases from uveal melanoma.²⁵⁻³³

Organ Procurement and Transplantation Network (OPTN). According to OPTN Policy, lesions eligible for down staging protocols to qualify for liver transplantation must meet one of the following criteria: One lesion greater than 5 cm and less than or equal to 8 cm or two or three lesions each greater than 3 cm and less than or equal to 5 cm, and a total diameter of all lesions less than or equal to 8 cm or four to five lesions each less than 3 cm, and a total diameter of all lesions less than or equal to 8 cm.⁴¹

American Association for the Study of Liver Disease (AASLD). *Practice Guidelines: Treatment of Hepatocellular Carcinoma* published in 2018 suggest that bridging to transplant with liver-directed therapy (LRT) in patients listed for liver transplantation within OPTN T2 (Milan) criteria to decrease progression of disease and subsequent dropout from the waiting list. The AASLD does not recommend one form of LRT over another for the purposes of bridging to liver transplantation for patients within OPTN T2 (Milan) criteria.³⁸

National Comprehensive Cancer Network (NCCN). *Clinical Practice Guidelines (CPG) in Oncology.*⁴⁰

- The CPG for hepatobiliary cancer (V1.2018), principles of locoregional therapy report that lesions 3 cm to 5 cm may be treated to prolong survival using arterially directed therapies, or with combination of an arterially directed therapy and ablation as long as tumor location is accessible for ablation. Unresectable/inoperable lesions >5 cm should be considered for treatment using arterially directed or systemic therapy. For bridge therapy to transplant arterially directed therapy is used to decrease tumor progression and the dropout rate from the liver transplant list and may be considered for patients who meet the transplant criteria.
- The CPG for neuroendocrine tumors of the gastrointestinal tract and/or distant metastases (V3.2017) includes a recommendation to consider hepatic-directed therapy for hepatic-predominant disease including arterial embolization and TACE for individuals with locoregional unresectable disease and/or distant (liver) metastases (symptomatic, clinically significant tumor burden, or clinically significant progressive disease).

Molina Clinical Policy
Transarterial Chemoembolization (TACE) and Transarterial Embolization (TAE) for Liver Tumors: Policy No. 120



Last Approval: 4/5/2021
 Next Review Due By: April 2022

- The CPG for uveal melanoma (V1.2018) includes the following recommendation: In general, uveal melanomas may have lower response rates to drug-based therapies than cutaneous melanoma, but efficacy has in general been more limited; however, individual patients on occasion may derive substantial benefit. Regionally directed therapies such as hepatic chemoembolization or radioembolization should be considered.
- The CPG for colorectal cancer (V2.2018) states that the data is not strong enough to recommend TACE for the treatment of colorectal liver metastases except in the setting of a clinical trial. The panel believes that arterial directed catheter therapy (radioembolization) is an option in highly selected patients with chemotherapy resistant refractive disease with predominant hepatic metastases.

SUPPLEMENTAL INFORMATION

None.

CODING & BILLING INFORMATION

CPT Codes

CPT	Description
37243	Vascular embolization or occlusion, inclusive of all radiological supervision and interpretation, intraprocedural roadmapping, and imaging guidance necessary to complete the intervention; for tumors, organ ischemia, or infarction
75894	Transcatheter therapy, embolization, any method, radiological supervision and interpretation

HCPCS Codes – N/A

ICD-10 Codes

ICD-10	Description
C22.0	Carcinoma malignant, hepatocellular
C22.9	Malignant neoplasm of liver, not specified as primary or secondary
C78.7	Secondary malignant neoplasm of liver and intrahepatic bile duct
C7A.00-C7A.8	Malignant neuroendocrine tumors
C7B.02	Secondary carcinoid tumors of liver
C69.30-C69.32	Malignant neoplasm of choroid
C69.40-C69.42	Malignant neoplasm of ciliary body

CODING DISCLAIMER. Codes listed in this policy are for reference purposes only and may not be all-inclusive. Deleted codes and codes which are not effective at the time the service is rendered may not be eligible for reimbursement. Listing of a service or device code in this policy does not guarantee coverage. Coverage is determined by the benefit document. Molina adheres to Current Procedural Terminology (CPT®), a registered trademark of the American Medical Association (AMA). All CPT codes and descriptions are copyrighted by the AMA; this information is included for informational purposes only. Providers and facilities are expected to utilize industry standard coding practices for all submissions. When improper billing and coding is not followed, Molina has the right to reject/deny the claim and recover claim payment(s). Due to changing industry practices, Molina reserves the right to revise this policy as needed.

APPROVAL HISTORY

- 9/18/2019, 4/23/2020, 4/5/2021** Policy reviewed, no changes, updated references.
- 7/10/2018** Policy reviewed and updated with revisions to criteria. For TACE and the addition of TAE for conditions including metastatic colorectal cancer, neuroendocrine tumors, uveal melanoma, as a bridge to liver transplant and in individuals who may become eligible for liver transplantation. Updated contraindications to TACE with additional recommendations Updated sections for General Information, Summary of Medical Evidence, Coding and References.
- 6/22/2017** Policy reviewed, no changes.
- 12/14/2016** Policy reviewed, no changes.
- 07/16/2015** Policy reviewed and updated with revisions to criteria (TACE utilizing chemotherapy-loaded microspheres [e.g., drug-loaded microspheres, drug-eluting beads, and doxorubicin drug-eluting bead transarterial chemoembolization (DEB-TACE)]; added Embosphere Microspheres are considered experimental, investigational and unproven for all liver-related conditions.
- 10/31/2012** New policy.

Molina Clinical Policy

Transarterial Chemoembolization (TACE) and Transarterial Embolization (TAE) for Liver Tumors: Policy No. 120

Last Approval: 4/5/2021

Next Review Due By: April 2022



REFERENCES

Government Agency

1. Centers for Medicare and Medicaid Services (CMS). Medicare coverage database. <http://www.cms.gov/mcd/search.asp>.

Peer Reviewed Publications

2. Chua TC, Liau W, Saxena A, Chu F, Glenn D, Chai A, Morris DL. Systematic review of neoadjuvant transarterial chemoembolization for resectable hepatocellular carcinoma. *Liver International* 2010; 30(2): 166-174. <http://onlinelibrary.wiley.com/doi/10.1111/j.1478-3231.2009.02166.x/full>.
3. Gruber-Rouh T, Naguib NN, Eichler K, Ackermann H, Zangos S, Trojan J, et al. Transarterial chemoembolization of unresectable systemic chemotherapy-refractory liver metastases from colorectal cancer: long-term results over a 10-year period. *Int J Cancer*. 2014 Mar 1;134(5):1225-31.
4. Llovet J, Bruix J: Systematic review of randomized trials for unresectable hepatocellular carcinoma: Chemoembolization improves survival. *Hepatology* 2003; 37:429-442.
5. Llovet J, Real M, Montanya X, et al: Arterial embolization, chemoembolization versus symptomatic treatment in patients with unresectable hepatocellular carcinoma: A randomized controlled trial. *Lancet* 2002; 359:1734-1739. [http://www.thelancet.com/journals/lancet/article/PIIS0140-6736\(02\)08649-X/fulltext](http://www.thelancet.com/journals/lancet/article/PIIS0140-6736(02)08649-X/fulltext)
6. Lo CM, Ngan H, Tso WK, et al: Randomized controlled trial of transarterial lipiodol chemoembolization for unresectable hepatocellular carcinoma. *Hepatology* 2002; 35:1164-1171. <http://onlinelibrary.wiley.com/doi/10.1053/jhep.2002.33156/abstract>.
7. Marelli L, Stigliano R, Triantos C, et al. Transarterial therapy for hepatocellular carcinoma: Which technique is more effective? A systematic review of cohort and randomized studies. *Cardiovasc Intervent Radiol*. 2007;30(1):6-25.
8. Morimoto M, Numata K, Kondou M, et al. Midterm outcomes in patients with intermediate-sized hepatocellular carcinoma: a randomized controlled trial for determining the efficacy of radiofrequency ablation combined with transcatheter arterial chemoembolization. *Cancer*. 2010 Dec 1;116(23):5452-60. doi: 10.1002/cncr.25314.
9. Oliveri RS, Wetterslev J, Gluud C. Transarterial (chemo) embolisation for unresectable hepatocellular carcinoma. *Cochrane Database of Systematic Reviews* 2011, Issue 3. Art. No.: CD004787. doi: 10.1002/14651858.CD004787.pub2
10. Samuel M, Chow PKH, Chan Shih-Yen E, Machin D, Soo K-C. Neoadjuvant and adjuvant therapy for surgical resection of hepatocellular carcinoma [update of *Cochrane Database Syst Rev* 2000; (2): CD001199; PMID: 10796754]. *Cochrane Database Syst Rev* 2009; (1): CD001199.
11. Takayasu K. Transarterial chemoembolization for hepatocellular carcinoma over three decades: current progress and perspective. *Jpn J Clin Oncol*. 2012 Apr;42(4):247-55. <http://www.ncbi.nlm.nih.gov/pubmed/22407946>.
12. Wang N, Guan Q, et al. TACE combined with PEI versus TACE alone in the treatment of HCC: A meta-analysis. *Med Oncol*. 2011 Dec;28(4):1038-43.
13. Zhang Z, Liu Q, He J, et al. The effect of preoperative transcatheter hepatic arterial chemoembolization on disease-free survival after hepatectomy for hepatocellular carcinoma. *Cancer* 2000; 89: 2606–12.
14. Zhong JH, Li LQ. Postoperative adjuvant transarterial chemoembolization for participants with hepatocellular carcinoma: A meta-analysis. *Hepatol Res*. 2010;40(10):943-953.
15. Zhou WP, Lai EC, Li AJ, et al. A prospective, randomized, controlled trial of preoperative transarterial chemoembolization for resectable large hepatocellular carcinoma. *Ann Surg* 2009; 249: 195–202.
16. Lammer J, Malagari K, Vogl T, et al. PRECISION V investigators: Prospective randomized study of doxorubicin-eluting-bead embolization in the treatment of hepatocellular carcinoma: results of the PRECISION V study. *Cardiovasc Intervent Radiol*. 2010;33(1):41-52. <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2816794/>.
17. Malagari K, Pomoni M, Kelekis A, et al. Prospective randomized comparison of chemoembolization with doxorubicin-eluting beads and bland embolization with BeadBlock for hepatocellular carcinoma. *Cardiovasc Intervent Radiol*. 2010;33(3):541-551.
18. Martin RC 2nd, Rustein L, Pérez Enguix D, et al. Hepatic arterial infusion of doxorubicin-loaded microsphere for treatment of hepatocellular cancer: a multi-institutional registry. *J Am Coll Surg*. 2011;213(4):493-500.
19. Dhanasekaran R, Kooby DA, Staley CA, Kauh JS, Khanna V, Kim HS. Comparison of conventional transarterial chemoembolization (TACE) and chemoembolization with doxorubicin drug eluting beads (DEB) for unresectable hepatocellular carcinoma (HCC). *J Surg Oncol*. 2010;101(6):476-480.
20. Song MJ, Park CH, Kim JD, et al. Drug-eluting bead loaded with doxorubicin versus conventional Lipiodol-based transarterial chemoembolization in the treatment of hepatocellular carcinoma: a case-control study of Asian patients. *Eur J Gastroenterol Hepatol*. 2011;23(6):521-527.
21. Malagari K, Alexopoulou E, Chatzimichael K, et al. Transcatheter chemoembolization in the treatment of HCC in patients not eligible for curative treatments: midterm results of doxorubicin-loaded DC bead. *Abdom Imaging*. 2008a;33(5):512-519.
22. Malagari K, Chatzimichael K, Alexopoulou E, et al. Transarterial chemoembolization of unresectable hepatocellular carcinoma with drug eluting beads: results of an open-label study of 62 patients. *Cardiovasc Intervent Radiol*. 2008b;31(2):269-280.
23. van Malenstein H, Maleux G, Vandecaveye V, et al. A randomized phase II study of drug-eluting beads versus transarterial chemoembolization for unresectable hepatocellular carcinoma. *Onkologie*. 2011;34(7):368-376.
24. Katsanos K, Kitrou P, et al. Comparative effectiveness of different transarterial embolization therapies alone or in combination with local ablative or adjuvant systemic treatments for unresectable hepatocellular carcinoma: A network meta-analysis of randomized controlled trials. *PLoS One*. 2017 Sep 21;12(9):e0184597. doi: 10.1371/journal.pone.0184597.
25. Grozinsky-Glasberg S, Kaltsas G, et al. Hepatic intra-arterial therapies in metastatic neuroendocrine tumors: lessons from clinical practice. *Endocrine*. 2018 Jun;60(3):499-509. doi: 10.1007/s12020-018-1537-0.
26. de Mestier L, Zappa M, et al. Liver transarterial embolizations in metastatic neuroendocrine tumors. *Rev Endocr Metab Disord*. 2017 Dec;18(4):459-471. doi: 10.1007/s11154-017-9431-2.

Molina Clinical Policy

Transarterial Chemoembolization (TACE) and Transarterial Embolization (TAE) for Liver Tumors: Policy No. 120

Last Approval: 4/5/2021

Next Review Due By: April 2022



27. Kennedy A, Bester L, et al. Role of hepatic intra-arterial therapies in metastatic neuroendocrine tumours (NET): guidelines from the NET-Liver-Metastases Consensus Conference. HPB (Oxford). 2015 Jan;17(1):29-37. doi: 10.1111/hpb.12326.
28. Do Minh D, Chapiro J, et al. Intra-arterial therapy of neuroendocrine tumour liver metastases: comparing conventional TACE, drug-eluting beads TACE and yttrium-90 radioembolisation as treatment options using a propensity score analysis model. Eur Radiol. 2017 Dec;27(12):4995-5005. doi: 10.1007/s00330-017-4856-2.
29. Valpione S, Aliberti C, Parrozzani R, et al. A retrospective analysis of 141 patients with liver metastases from uveal melanoma: a two-cohort study comparing transarterial chemoembolization with CPT-11 charged microbeads and historical treatments. Melanoma Res. 2015 Apr;25(2):164-8. doi: 10.1097/CMR.000000000000129.
30. Duran R, Chapiro J et al. Uveal melanoma metastatic to the liver: The role of quantitative volumetric contrast-enhanced mr imaging in the assessment of early tumor response after transarterial chemoembolization. Transl Oncol. 2014 Aug;7(4):447-55. doi: 10.1016/j.tranon.2014.05.004.
31. Shibayama Y, Namikawa K, et al. Efficacy and toxicity of transarterial chemoembolization therapy using cisplatin and gelatin sponge in patients with liver metastases from uveal melanoma in an Asian population. Int J Clin Oncol. 2017 Jun;22(3):577-584. doi: 10.1007/s10147-017-1095-0.
32. Riemsma RP, Bala MM, Wolff R, Kleijnen J. Transarterial (chemo)embolization versus no intervention or placebo intervention for liver metastases. Cochrane Database Syst Rev. 2013;(4):CD009498.
33. Belinson S, Chopra R, Yang Y, et al. Local hepatic therapies for metastases to the liver from unresectable colorectal cancer: Comparative effectiveness review (no. 93). Publication No. 13-EHC014-EF. Agency for Healthcare Research and Quality (AHRQ). December 2012. Archived. <https://effectivehealthcare.ahrq.gov/topics/colorectal-cancer-liver-metastases-therapies/research>.
34. Hayes. <https://evidence.hayesinc.com>. Registration and login required.
 - Health technology assessment: Doxorubicin drug-eluting bead transarterial chemoembolization (DEB-TACE) for patients with hepatocellular carcinoma (HCC). Published April 2014. Archived 2015.
 - Health technology assessment: Transarterial embolization with embozene microspheres (CeloNova BioSciences Inc.) for Treatment of Hepatocellular Carcinoma and Other Liver Tumors. Updated August 2015. Archived 2016.
35. Hayes. Health technology assessment: Transarterial chemoembolization plus radiofrequency ablation for liver cancer. <https://evidence.hayesinc.com>. Published 2012. Updated 2016. Archived 2018. Registration and login required.

National and Specialty Organizations

36. American College of Radiology (ACR). ACR appropriateness criteria radiologic management of hepatic malignancy. Updated 2015. <https://acsearch.acr.org/docs/69379/Narrative/>.
37. Gaba RC, Lokken P, Hickey RM, et al. Society of Interventional Radiology: Quality improvement guidelines for transarterial chemoembolization and embolization of hepatic malignancy. 2017; (9):1210-1223.
38. American Association for the Study of Liver Disease (AASLD). Clinical practice guidelines. <http://www.aasld.org/publications/practice-guidelines-0>.
 - Heimbach, et al. Treatment of hepatocellular carcinoma. Hepatology. 2018. 67(1):358-380.
 - Martin P, et al. Liver transplantation: Evaluation of the adult patient. Hepatology. 2014 Mar;59(3):1144-65.
39. National Cancer Institute (NCI).
 - Childhood liver cancer treatment PDQ®. Updated 2021. <https://www.cancer.gov/types/liver/hp/child-liver-treatment-pdq>.
 - Adult primary liver cancer treatment PDQ®. Updated 2021. <https://www.cancer.gov/types/liver/hp/adult-liver-treatment-pdq>.
40. National Comprehensive Cancer Network (NCCN). Clinical practice guidelines in oncology (hepatobiliary cancers; neuroendocrine tumors of the gastrointestinal tract and/or distant metastases; uveal melanoma; colon cancer). <http://www.nccn.org>.
41. Organ Procurement and Transplantation Network (OPTN). Organ distribution: Policy 9: allocation of livers and liver-intestines. Effective January 18, 2020. <https://optn.transplant.hrsa.gov/resources/by-organ/liver-intestine/>.

Other Evidence Based Reviews and Publications

42. UpToDate. <http://www.uptodate.com>. Registration and login required.
 - Curley SA, Stuart KE, Schwartz JM. Nonsurgical therapies for localized hepatocellular carcinoma: Transarterial embolization, radiotherapy, and radioembolization.
 - Carvajal R. Management of metastatic uveal melanoma.
43. DynaMed. EBSCO Publishing. 1995-2020. Hepatocellular carcinoma.
44. Advanced Medical Review (AMR).
 - AMR Peer Review. Policy reviewed on October 15, 2012 by an Advanced Medical Reviews (AMR) practicing, board-certified physician in the areas of Internal Medicine, Oncology, Hematology.
 - AMR Peer Review. Policy reviewed on July 31, 2018 by an Advanced Medical Reviews (AMR) practicing, board-certified physician in the area of Radiation Oncology.
45. McKesson InterQual. Procedures: Ablative or transarterial therapy, liver. Updated 2019.

APPENDIX

Reserved for State specific information (to be provided by the individual States, not Corporate). Information includes, but is not limited to, State contract language, Medicaid criteria and other mandated criteria.